

DUPIXENT[®]
(dupilumab) Injection
200mg • 300mg

**HELP
HEAL
YOUR
SKIN
FROM
WITHIN**[™]

STAY AHEAD OF ECZEMA WITH DUPIXENT

FDA-approved for ages 6 months and up with uncontrolled moderate-to-severe eczema (atopic dermatitis)



**SHOW OFF
TO THE WORLD**

Actual patient.
Individual results may vary.

INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 6 months of age and older with moderate-to-severe eczema (atopic dermatitis or AD) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids. It is not known if DUPIXENT is safe and effective in children with atopic dermatitis under 6 months of age.

Please see Important Safety Information throughout and click [here](#) for full [Prescribing Information](#) and [Patient Information](#).


**CRISTAL,
ACTUAL PATIENT**

KNOW WHAT TO LOOK FOR

Uncontrolled moderate-to-severe eczema (atopic dermatitis) can be difficult to manage. You might do everything right and still experience symptoms. But have you considered all your options? You may want to ask your doctor about making a change to your treatment.

WHEN SHOULD YOU TALK TO AN ECZEMA SPECIALIST? WHEN YOU OR YOUR CHILD:

- ▶ HAVE ECZEMA THAT KEEPS COMING BACK
- ▶ TAKE ORAL STEROIDS MORE THAN ONCE A YEAR
- ▶ TAKE IMMUNOSUPPRESSANTS MORE THAN ONCE A YEAR
- ▶ HIDE SKIN FROM OTHERS



Remember that good communication is the first step towards feeling better about your eczema. Scan code or [CLICK HERE](#) to download our Doctor Discussion Guide and get this conversation started with your specialist.

DID YOU KNOW?

- ▶ YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®. Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you: have eye problems; have a parasitic (helminth) infection; are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT. Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

A DIFFERENCE INSIDE AND OUT

When topical prescriptions aren't enough, and your symptoms and flare-ups keep coming back, it might be time for a different kind of treatment like a biologic. And DUPIXENT is just that.

DUPIXENT is a breakthrough treatment for uncontrolled moderate-to-severe eczema.

DUPIXENT:

- ▶ IS A BIOLOGIC THAT HELPS BLOCK A KEY SOURCE OF INFLAMMATION INSIDE THE BODY THAT CAN CAUSE ECZEMA
- ▶ HELPS YOU STAY AHEAD OF SYMPTOMS FOR CLEARER SKIN AND LESS ITCH
- ▶ CONTINUOUSLY TREATS ECZEMA, EVEN BETWEEN FLARE-UPS
- ▶ IS NOT A STEROID OR IMMUNOSUPPRESSANT
- ▶ IS NOT TAKEN DAILY



**BECKETT,
ACTUAL PATIENT**

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you (cont'd): are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby. A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>; are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**ADANTE,
ACTUAL PATIENT**

**OVER
457,000*
PATIENTS
PRESCRIBED DUPIXENT**

*New adult and pediatric patients (6 months of age and older) with uncontrolled moderate-to-severe eczema.

IQVIA National Source of Business (NSOB) data through December 2023



**ANTHONY,
ACTUAL PATIENT**



SCAN CODE OR [CLICK HERE](#)
TO LEARN MORE
ABOUT HOW
DUPIXENT CAN HELP

TAKE CONTROL TAKE YOUR SHOT AGAINST ECZEMA

Many people with moderate-to-severe eczema have an overly active immune system that produces more inflammation under the skin than usual.

DUPIXENT helps block a key source of inflammation that may be causing your eczema symptoms. It works continuously on the inside to help heal your eczema on the outside. And, you don't take it every day—who wants to be reminded of their eczema on a daily basis?

DUPIXENT is a biologic—biologics are specialty medicines used to treat a variety of conditions, including chronic inflammatory diseases such as moderate-to-severe eczema. Biologics are processed in the body differently than oral or topical medications. To be effective and work in the body properly, most biologics are injectable medicines due to their size and the unique way they are made.



**JOLIE,
ACTUAL PATIENT**

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

MEANINGFUL RESULTS FROM CLINICAL TRIALS



HOW WERE THE TRIALS SET UP?

DUPIXENT was well studied in 6 clinical trials with 2,899 patients, ages 6 months through adult, to assess its safety profile, and to determine its efficacy in clearing skin and reducing itch.

Every age group studied had moderate and/or severe eczema that was uncontrolled with topical prescription medications.

DUPIXENT was first approved to treat adults with uncontrolled moderate-to-severe eczema by the FDA in 2017.



WHAT DID THE TRIALS MEASURE?

The trials measured skin clearance, itch reduction, and safety among people taking DUPIXENT compared to those who were not taking DUPIXENT. Each trial measured the results at 16 weeks. One trial measured the results at 16 and 52 weeks (1 year).



WHAT MEDICATIONS WERE USED?

Depending on the trial, participants received one of the following treatments:

- ▶ DUPIXENT
- ▶ Placebo (a treatment that appears to be real but contains no medicine)
- ▶ DUPIXENT with a topical corticosteroid (TCS)
- ▶ Placebo with TCS

DID YOU KNOW?

- ▶ YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**KATHY,
ACTUAL PATIENT**

ADANTE,
ACTUAL PATIENT

~5 YEAR SAFETY RESULTS IN ADULTS...



LONG-TERM SAFETY RESULTS FOR ADULTS IN AN EXTENSION STUDY WERE CONSISTENT WITH THOSE SEEN IN THE ORIGINAL CLINICAL TRIALS

The most common side effects observed in clinical trials were: injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, dry eye, cold sores in the mouth or on the lips, and high count of a certain white blood cell (eosinophilia).

...AND IMPORTANT CONSIDERATIONS



STUDIED IN 15 CLINICAL TRIALS
(6 IN ECZEMA, 9 IN 4 OTHER CONDITIONS)



#1 PRESCRIBED BIOLOGIC FOR ECZEMA BY
DERMATOLOGISTS AND ALLERGISTS



NOT A STEROID OR IMMUNOSUPPRESSANT



NO INITIAL OR ROUTINE BLOOD WORK
REQUIRED, PER PRESCRIBING INFORMATION



CAN BE USED WITH OR WITHOUT
TOPICAL CORTICOSTEROIDS

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

ANTHONY,
ACTUAL PATIENT

SHOW OFF CLEARER SKIN AND LESS ITCH

TRIAL RESULTS: DUPIXENT vs PLACEBO

TWO SEPARATE 16-WEEK CLINICAL TRIALS SHOWED:

~**4X** SAW CLEAR OR ALMOST CLEAR SKIN

37% vs 9% not taking DUPIXENT (placebo)

>**3X** HAD SIGNIFICANT ITCH REDUCTION

38% vs 11% not taking DUPIXENT

ALMOST HALF OF ADULTS SAW

≥**75%** CLEARER SKIN

48% vs 13% not taking DUPIXENT

MANY SAW

≥**90%** CLEARER SKIN

33% vs 7% not taking DUPIXENT

IMPORTANT SAFETY INFORMATION: Especially tell your healthcare provider if you are taking oral, topical or inhaled corticosteroid medicines or if you have atopic dermatitis and asthma and use an asthma medicine.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



SCAN CODE OR CLICK HERE
TO HEAR ACTUAL
PATIENT STORIES.

ADULTS (AGES 18+)

SHOW OFF WHAT'S POSSIBLE IN THE LONG RUN

TRIAL RESULTS: DUPIXENT + TOPICAL CORTICOSTEROIDS (TCS) vs TCS ONLY

LONG-TERM RESULTS FROM A CLINICAL TRIAL
THAT STUDIED DUPIXENT FOR 52 WEEKS SHOWED:

CLEAR OR ALMOST CLEAR SKIN AT 16 WEEKS

39% taking DUPIXENT + TCS vs 12% using TCS only

LONG-LASTING CLEARER SKIN AT 16 AND 52 WEEKS

22% taking DUPIXENT + TCS vs 7% using TCS only

FAST ITCH RELIEF AS SOON AS 2 WEEKS

18% taking DUPIXENT + TCS vs 8% using TCS only

SIGNIFICANT ITCH REDUCTION AT 16 WEEKS

59% taking DUPIXENT + TCS vs 20% using TCS only

ITCH RELIEF THAT LASTS AT 52 WEEKS

51% taking DUPIXENT + TCS vs 13% using TCS only

IMPORTANT SAFETY INFORMATION: Do not change or stop your corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine or other asthma medicine to come back.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**ADANTE,
ACTUAL PATIENT**

STAY AHEAD WITH CLEARER SKIN AND LESS ITCH

TRIAL RESULTS: DUPIXENT vs PLACEBO

TEENS TAKING PART IN A 16-WEEK CLINICAL TRIAL IN WHICH:

12X SAW CLEAR OR ALMOST CLEAR SKIN

24% vs 2% not taking DUPIXENT (placebo)

~7X EXPERIENCED ITCH RELIEF

37% vs 5% not taking DUPIXENT

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including: Allergic reactions. DUPIXENT can cause allergic reactions that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

TYLER,
ACTUAL PATIENT

TEENS (AGES 12-17)

PROVEN RESULTS AND DEMONSTRATED SAFETY PROFILE IN TEENS

TRIAL RESULTS CONTINUED: DUPIXENT vs PLACEBO

TEENS TOOK PART IN A 16-WEEK CLINICAL TRIAL IN WHICH:

NEARLY HALF OF TEENS SAW
≥75% CLEARER SKIN

42% vs 8% in those not taking DUPIXENT

SOME EVEN SAW
≥90% CLEARER SKIN

23% vs 2% not taking DUPIXENT

► **YOU CAN TAKE DUPIXENT WITH OR WITHOUT
A TOPICAL CORTICOSTEROID (TCS)**

SAFETY RESULTS:

Long-term safety results for teens continuing on DUPIXENT in the pediatric extension study through 1 year were consistent with the adult clinical trials.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): Eye problems. Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**JOLIE,
ACTUAL PATIENT**

RESULTS THEY CAN FEEL RESULTS THEY CAN SHOW OFF

Caring for your child with uncontrolled moderate-to-severe eczema can be a struggle—especially when you're constantly trying to avoid triggers.

TRIAL RESULTS: DUPIXENT + TCS vs TCS ONLY

CHILDREN TAKING PART IN A 16-WEEK CLINICAL TRIAL IN WHICH:

>2X SAW CLEAR OR ALMOST CLEAR SKIN

30% on 300 mg every 4 weeks + TCS vs 13% on TCS only
39% on 200 mg every 2 weeks + TCS vs 10% on TCS only

~4.5X HAD LESS ITCH

54% on 300 mg every 4 weeks + TCS vs 12% on TCS only
61% on 200 mg every 2 weeks + TCS vs 13% on TCS only

YOUR HEALTHCARE PROVIDER WILL TELL YOU HOW MUCH DUPIXENT TO INJECT AND HOW OFTEN

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): Eye problems. Your healthcare provider may send you to an ophthalmologist for an eye exam if needed.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

CALLA,
ACTUAL PATIENT

CHILDREN (AGES 6-11)

PROVEN RESULTS AND DEMONSTRATED SAFETY PROFILE IN CHILDREN

TRIAL RESULTS CONTINUED: DUPIXENT + TCS vs TCS ONLY

CHILDREN TOOK PART IN A 16-WEEK CLINICAL TRIAL IN WHICH:

MORE THAN HALF OF CHILDREN SAW

≥75% CLEARER SKIN

75% on 300 mg every 4 weeks + TCS vs 28% on TCS only

75% on 200 mg every 2 weeks + TCS vs 26% on TCS only

YOUR HEALTHCARE PROVIDER WILL TELL YOU HOW MUCH DUPIXENT TO INJECT
AND HOW OFTEN

► YOU CAN TAKE DUPIXENT WITH OR WITHOUT
A TOPICAL CORTICOSTEROID (TCS)

SAFETY RESULTS:

Long-term safety results for children continuing on DUPIXENT with or without TCS in the pediatric extension study through 1 year were consistent with the adult clinical trials.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): Joint aches and pain. Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

BECKETT,
ACTUAL PATIENT



**CAMERON,
ACTUAL PATIENT**

HEALING FROM WITHIN FOR YOUR LITTLE ONE

DUPIXENT is the only FDA-approved biologic for treating uncontrolled moderate-to-severe eczema in children as young as 6 months.

TRIAL RESULTS: DUPIXENT + TCS vs TCS ONLY

YOUNG CHILDREN TAKING PART IN A 16-WEEK CLINICAL TRIAL IN WHICH:

7X SAW CLEAR OR ALMOST CLEAR SKIN

28% on DUPIXENT + TCS vs 4% on TCS only

~5X HAD LESS ITCH

48% on DUPIXENT + TCS vs 9% on TCS only

HALF OF YOUNG CHILDREN SAW

≥75% CLEARER SKIN

53% on DUPIXENT + TCS vs 11% on TCS only

▶ **YOU CAN TAKE DUPIXENT WITH OR WITHOUT
A TOPICAL CORTICOSTEROID (TCS)**

SAFETY RESULTS:

Long-term safety results in young children continuing on DUPIXENT with or without TCS in the pediatric extension study through 1 year were consistent with the adult clinical trials. In addition, hand-foot-and-mouth disease and warts were reported. In these reported cases, patients continued treatment with DUPIXENT.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): Joint aches and pain. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

HELPS HEAL THE LOOK AND FEEL OF SKIN

Eczema can affect anyone, regardless of age, ethnicity, or gender, and it can appear differently on different skin types. In some cases, eczema rashes may look red, gray, brown, or purple. DUPIXENT has shown generally consistent efficacy regardless of race/ethnicity in adult studies.



Actual 4-year-old patient in the clinical trial.[†] Individual results may vary.



Actual adult patient.*



Actual 6-year-old patient in the clinical trial.[†] Individual results may vary.

*This adult patient was an actual patient treated with DUPIXENT. Not clinical trial patient. Scoring was designated by the treating physician. Because this was a real-world patient, other factors may have influenced their treatment results. Individual results may vary.

[†]All patients were prescribed low or medium-potency topical corticosteroids (TCS) based on the clinical trial program.

IMPORTANT SAFETY INFORMATION: The most common side effects in patients with eczema include injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, dry eye, cold sores in your mouth or on your lips, and high count of a certain white blood cell (eosinophilia). Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

DOSING FOR DUPIXENT® (dupilumab):

	BODY WEIGHT	INITIAL LOADING DOSE	SUBSEQUENT DOSAGE
ADULTS 18+	Any weight	600 mg (two 300-mg injections)	300 mg; every 2 weeks
CHILDREN & TEENS 6 TO 17 YEARS OF AGE	132 lb or more	600 mg (two 300-mg injections)	300 mg; every 2 weeks
	66 lb to under 132 lb	400 mg (two 200-mg injections)	200 mg; every 2 weeks
	33 lb to under 66 lb	600 mg (two 300-mg injections)	300 mg; every 4 weeks
	BODY WEIGHT	INITIAL* AND SUBSEQUENT DOSAGE	
YOUNG CHILDREN 6 MONTHS TO 5 YEARS OF AGE	33 lb to under 66 lb	300 mg; every 4 weeks	
	11 lb to under 33 lb	200 mg; every 4 weeks	

*No loading dose recommended for young children.

DUPIXENT is available as a pre-filled pen or pre-filled syringe.

Your doctor will tell you how much DUPIXENT to inject and how often to inject it, following proper training. DUPIXENT is an injection given under the skin (subcutaneous injection). Please rotate the injection site every time you administer DUPIXENT.


DUPIXENT can be given by a doctor in the office or a caregiver at home:

- ▶ in children 6 months to less than 12 years, DUPIXENT should be given by a caregiver
- ▶ in teens 12–17 years, DUPIXENT should be given by or supervised by an adult

YOUR HEALTHCARE PROVIDER WILL DECIDE IF YOU OR YOUR CAREGIVER CAN INJECT DUPIXENT.



Pre-filled pen: available in 200 mg and 300 mg doses for patients aged 2+ years



SCAN CODE OR [CLICK HERE](#) TO WATCH THE SUPPLEMENTAL TRAINING VIDEOS ON HOW TO ADMINISTER DUPIXENT



Pre-filled syringe: available in 200 mg and 300 mg doses for patients aged 6+ months

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088. Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

DUPIXENT AND YOU

A PERSONALIZED DOCTOR DISCUSSION GUIDE

Whether you're just beginning your treatment journey or looking for another way to treat your uncontrolled moderate-to-severe eczema, the following checklist is for both patients and caregivers. Think of this guide as the start of a conversation you can continue with your eczema specialist to help determine if DUPIXENT may be right for you or your loved one. Choose all that apply.

Are you a patient or a caregiver?

- Patient Caregiver

What are your or your loved one's treatment goals?

- Clearer skin Itch relief

How long have you or your loved one had moderate-to-severe eczema?

- 0-2 years 2-5 years 5-10 years 10+ years

What treatments have been prescribed or used for your or your loved one's eczema?

- Topical steroids Biologics None
 Phototherapy Over-the-counter products
 Oral steroids Other _____

Describe your or your loved one's eczema symptoms.

RASH SEVERITY

- None Mild Moderate Severe

ITCH INTENSITY

- None Mild Moderate Severe

IMPORTANT SAFETY INFORMATION: Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

How often do these symptoms occur?

ITCHING

- Never Rarely Occasionally Frequently

INFLAMED, DISCOLORED SKIN

- Never Rarely Occasionally Frequently

THICKENED, HARDENED PATCHES OF SKIN

- Never Rarely Occasionally Frequently

How often do flare-ups occur...

- Every few months Monthly Weekly Daily

...and for how long?

- 1-5 days 5-10 days 10-15 days 15+ days

Where are your or your loved one's eczema symptoms?

- Neck Chest Stomach Front of Knees
 Face Elbows Feet Back of Knees
 Hands Ankles Scalp Other _____

Describe any additional impact living with eczema has had on you or your loved one.



Now that you've finished, just snap a photo to share with your eczema specialist at your next appointment

DUPIXENT myway

WITH YOU EVERY STEP OF THE WAY

Once you've been prescribed DUPIXENT, the *DUPIXENT MyWay*® patient support program is here to help answer questions you may have about DUPIXENT. Keep in mind your doctor is still the best resource for you if you have any questions about your eczema or your treatment program.

To enroll in *DUPIXENT MyWay*® and access helpful resources, call 1-844-DUPIXENT (1-844-387-4936) and select Option 1.



CAMERON,
ACTUAL PATIENT

YOU'RE NOT ALONE

It's important to know that *DUPIXENT MyWay*® helps adults and children 6 months and up who are on or about to start DUPIXENT.



Once enrolled in the program, and while your insurance benefits are being confirmed for DUPIXENT, you will receive a Welcome Call from your **Case Manager**, who will explain the support available from *DUPIXENT MyWay*. In addition, your **Case Manager** can offer education and support to help you start and stay on track with your treatment.




Your *DUPIXENT MyWay Case Manager* can answer questions about DUPIXENT, arrange supplemental injection training, and help you start and stay on track.



With the *DUPIXENT MyWay Patient App*, people just like you can get access to tools to help you start and stay on track with your treatment. Contact your **Case Manager** for an invitation with directions on how to get the app.



The *DUPIXENT MyWay Mentor Program* gives you the opportunity to speak with other DUPIXENT patients and caregivers to hear their experiences.



SCAN CODE OR [CLICK HERE](#) TO LEARN ABOUT ADDITIONAL DUPIXENT MYWAY RESOURCES.

IMPORTANT SAFETY INFORMATION: Do not try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 months to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

DUPIXENT myway

NEED HELP PAYING?

The DUPIXENT MyWay Copay Card Program may help reduce the out-of-pocket cost of DUPIXENT. Eligible patients may pay as little as \$0* per fill for each DUPIXENT prescription (maximum of \$13,000 per patient per calendar year)*.

AS LITTLE AS \$0* COPAY MAY BE AVAILABLE IF YOU:

- ▶ Have commercial insurance
- ▶ Have a DUPIXENT prescription for an FDA-approved condition
- ▶ Reside in the 50 United States, the District of Columbia, Puerto Rico, Guam, or the USVI
- ▶ Are a patient or caregiver 18 years or older

**Approval is not guaranteed. Program has an annual maximum of \$13,000. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed or restricted. DUPIXENT MyWay reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.*

OTHER OPTIONS FOR SUPPORT

Through the Patient Assistance Program, eligible patients who are uninsured or rendered uninsured by their insurance provider could receive DUPIXENT free of charge if they qualify.

Those who may qualify must be at least 18 years of age or older, a resident of the 50 United States, the District of Columbia, Puerto Rico, Guam, or the USVI, and demonstrate a financial need.

Call DUPIXENT MyWay® at 1-844-387-4936 for full program information and eligibility criteria.



**KATHY,
ACTUAL PATIENT**

DUPIXENT 
(dupilumab) Injection
200mg · 300mg

HELP HEAL YOUR SKIN FROM WITHIN™

ASK YOUR ECZEMA SPECIALIST
ABOUT DUPIXENT AND
VISIT [DUPIXENT.COM](https://www.dupixent.com)
OR CALL 1-844-DUPIXENT
(1-844-387-4936)



SCAN CODE OR
[CLICK HERE](#)
TO LEARN MORE
ABOUT DUPIXENT



IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in **DUPIXENT**®.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

sanofi | **REGENERON**®

© 2024 Sanofi and Regeneron Pharmaceuticals, Inc. All rights reserved.
DUPIXENT® and DUPIXENT MyWay® are registered trademarks of
Sanofi Biotechnology. US.DUP.23.09.0173 02/24