



DU MORE WITH LESS ASTHMA

#1 PRESCRIBED BIOLOGIC BY PULMONOLOGISTS AND ALLERGISTS

In an aggregate across all indications. Data on file, Sanofi US. IQVIA weekly NSOB; data through June 2023.

APPROVED FOR AGES 6 AND UP

INDICATION: DUPIXENT is a prescription medicine used with other asthma medicines for the maintenance treatment of moderate-to-severe eosinophilic or oral steroid dependent asthma in adults and children 6 years of age and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing. DUPIXENT is not used to treat sudden breathing problems. It is not known if DUPIXENT is safe and effective in children with asthma under 6 years of age.

Please see additional **Important Safety Information** throughout and accompanying full **Prescribing Information** including **Patient Information**.

DUPIXENT 
(dupilumab) Injection
200mg · 300mg

THE NEXT STEPS

TALK TO YOUR DOCTOR

THE RESULTS

THE SIGNS

THE BENEFITS



BENEFITS WITH EVERY BREATH



TARGETS A SOURCE OF ASTHMA DRIVEN BY TYPE 2 INFLAMMATION TO HELP IMPROVE SYMPTOMS



IMPROVES LUNG FUNCTION SO YOU OR YOUR CHILD CAN BREATHE BETTER IN AS LITTLE AS **2 WEEKS**



HELPS PREVENT ASTHMA ATTACKS



CAN REDUCE OR COMPLETELY ELIMINATE ORAL STEROID USE

Prescribed to over **100,000** asthma patients and counting!^{1,a,b}

^aLH Patient View; data through June 2023.

^bNew adult and pediatric patients aged 6 years and older with moderate-to-severe eosinophilic or OCS-dependent asthma.


DUPIXENT was studied in 4 clinical trials with more than 3,200 people 6 years of age and older with moderate-to-severe asthma.

IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

A FIRST-OF-ITS-KIND ASTHMA TREATMENT


Scientific advances have found that an underlying inflammation, known as **type 2 inflammation** may be a source of uncontrolled moderate-to-severe asthma.



DID YOU KNOW
8 OF 10
ASTHMA PATIENTS HAVE
TYPE 2 INFLAMMATION?

DUPIXENT blocks two of the key sources of type 2 inflammation in the lungs. This can result in a reduction of asthma symptoms.

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you: have a parasitic (helminth) infection; are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT.



IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you (cont'd): are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby. A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

THE NEXT STEPS

TALK TO YOUR DOCTOR

THE RESULTS

THE SIGNS

THE BENEFITS

BETTER BREATHING. LESS ORAL STEROIDS.

While oral steroids may be helpful when prescribed, there is some medical concern over long-term use of oral steroids, which includes:

- weight gain
- diabetes
- mood change
- vision problems

In a study of people dependent on oral corticosteroids:

86% OF PEOPLE ON DUPIXENT
REDUCED OR ELIMINATED
THEIR ORAL STEROID USE*†

*In a clinical study of patients 12+ years old with DUPIXENT 300 mg + standard of care (SOC) vs **68%** with placebo + SOC.

†Do not stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you (cont'd): are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DO YOU KNOW THE SIGNS OF CONTROL?

People often think their asthma or their child's asthma is under control, but it may not be. There may be signs that you or your child could be doing better.

SOME KEY FACTORS THAT MAY INDICATE THAT MODERATE-TO-SEVERE ASTHMA ISN'T WELL-CONTROLLED:



Rescue inhaler use is more than twice a week



In the past month, sleep has been interrupted because of asthma



Activities were limited or missed because of an attack



A trip to the ER because of asthma occurred at least once in the past year

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.



IMPORTANT SAFETY INFORMATION: Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or use an asthma medicine. **Do not** change or stop your corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine or other asthma medicine to come back.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DU MORE YOU

In clinical studies, people taking DUPIXENT saw a meaningful improvement in their day-to-day lives, which included:

- LESS SHORTNESS OF BREATH
- LESS WHEEZING
- NOT WAKING AT NIGHT
- BEING LESS LIMITED IN THEIR NORMAL ACTIVITIES

The relationship between DUPIXENT and these results were studied but not definitively proven.

The most common side effects include injection site reactions, pain in the throat (oropharyngeal pain), and high count of a certain white blood cell (eosinophilia).

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including: Allergic reactions. DUPIXENT can cause allergic reactions that can sometimes be severe.

Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.



DU LESS ASTHMA

EXPERIENCED
UP TO **81%** FEWER
ASTHMA ATTACKS*†

Clinical studies have shown that DUPIXENT can:

- HELP SIGNIFICANTLY IMPROVE LUNG FUNCTION
- HELP PREVENT SEVERE ASTHMA ATTACKS

*In those with high eosinophils (a type of white blood cell).

†The annual rate of severe asthma attacks was 0.20 with DUPIXENT 300 mg vs 1.04 with placebo.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): Inflammation of your blood vessels. Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. It is not known whether this is caused by DUPIXENT. Tell your healthcare provider right away if you have: rash, chest pain, worsening shortness of breath, a feeling of pins and needles or numbness of your arms or legs, or persistent fever.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): Joint aches and pain. Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

YOUR CHILD DESERVES TO DO MORE TOO

And with less asthma, they can. DUPIXENT is proven to improve lung function and help prevent asthma attacks in children 6-11 years of age.

65% FEWER ASTHMA ATTACKS
IN CHILDREN*

*The annual rate of severe asthma attacks was 0.24 with DUPIXENT 100/200 mg vs 0.67 with placebo.

66% REDUCTION IN STEROID
BURSTS IN CHILDREN†

†Do not stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.

IMPORTANT SAFETY INFORMATION: The most common side effects in patients with asthma include injection site reactions, high count of a certain white blood cell (eosinophilia), pain in the throat (oropharyngeal pain), and parasitic (helminth) infections.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.




REAL EXPERIENCES FROM REAL PEOPLE

When I was first diagnosed with asthma I was 23 years old. I had no previous symptoms, no family history, and I just started wheezing and coughing. Since I've started using DUPIXENT, I'm able to do more things that I love, like playing with my grandchildren.



Marlena
DUPIXENT MyWay®
Patient ambassador

I'm an outdoors guy. But I couldn't always be as active as I wanted to be. I was constantly worried about my asthma. Now, I'm able to DU MORE of the things I like to do. I'm chasing rainbow trout in the mountains. I'm feeding cattle on the farm.



Tim
DUPIXENT MyWay
Patient ambassador

I have been living with severe asthma for ten years. My symptoms and the coughing would bring on an asthma attack. DUPIXENT has worked for me. It helps me breathe.



Pamela
DUPIXENT MyWay
Patient ambassador

Individual results may vary.

Hearing from people who have had a similar experience can help.

To hear more stories of how DUPIXENT has helped others, click [here](#).



Join one of our webinars where you can learn from healthcare professionals and patients who know what you're going through.

Visit [DUPIXENT.com/as-webinars](https://www.dupilumab.com/as-webinars) to see what may be available.

IMPORTANT SAFETY INFORMATION: Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

A HELPFUL GUIDE

Be sure to bring this with you to your next appointment. A guided conversation can be the first step to getting better.

1. How long have you been experiencing asthma symptoms?

_____ Year(s) _____ Month(s)

2. What treatments have you or your loved one been prescribed for asthma?

Select all that apply:

- Inhalers Nebulizers Biologics
 Inhaled corticosteroids Inhaled long-acting beta antagonists
 Oral corticosteroids None Other _____

3. How often do these symptoms occur?

Coughing

- Never Rarely Occasionally Frequently

Wheezing

- Never Rarely Occasionally Frequently

Rapid Breathing

- Never Rarely Occasionally Frequently

Chest Tightness

- Never Rarely Occasionally Frequently

4. Have you woken up at night because of uncontrolled moderate-to-severe asthma in the past month?

- Yes No

5. Have you limited or missed an activity because of an asthma attack?

- Yes No

6. Have you used your rescue inhaler more than twice during a typical week?

- Yes No

7. Have you noticed any of the following triggers for your asthma?

- Dust mites Air pollution Pets Mold
 Perfumes Scented soaps Physical Exercise
 Other _____

HAS YOUR DOCTOR PRESCRIBED DUPIXENT[®] (dupilumab)?

Your healthcare provider is your best resource for information about DUPIXENT and should be your primary point of contact throughout your treatment journey.


YOU BEGIN YOUR JOURNEY.

Your healthcare provider prescribes DUPIXENT, and you enroll into *DUPIXENT MyWay*.

- Your healthcare provider can download the enrollment form on [DUPIXENT.com](https://www.dupixent.com), help you fill it out, and fax it back to *DUPIXENT MyWay* at 1-844-387-9370
- You or your healthcare provider can call 1-844-DUPIXEN(T) Option 1
- Providing your email address allows *DUPIXENT MyWay* to give you more support resources about DUPIXENT

QUESTIONS ABOUT INJECTING? RESOURCES ARE HERE FOR YOU.

Whether you've received injection training from your healthcare provider but are new to injecting or have questions about taking DUPIXENT, we're with you.



Take advantage of our supplemental injection resources and training videos by scanning the QR code or clicking [here](#).

WHAT CAN *DUPIXENT MyWay*[®] DO FOR YOU?

Enrolling in *DUPIXENT MyWay* can help ensure you receive DUPIXENT (dupilumab) as quickly as possible and receive additional support along your treatment journey. For eligible patients, *DUPIXENT MyWay* can:



Help research and explain your insurance benefits



Find ways to help with the out-of-pocket cost of DUPIXENT if you need help with your copay, you are denied coverage from your health plan, or you do not have or experience a loss in insurance



Connect you with a Nurse Educator to answer questions about DUPIXENT and help you start and stay on track with therapy



Connect you with your specialty pharmacy to make sure your prescription is shipped to your home or preferred location



Explain how to properly store DUPIXENT when you receive your shipment



Provide supplemental injection training—in person, virtually, or over the phone—to help you or a caregiver get more familiar with injecting DUPIXENT



Offer a needle disposing kit or sharps container for proper disposal



Remind you when it is time to refill your DUPIXENT prescription

NEED HELP PAYING? \$0* COPAY MAY BE AVAILABLE

Eligible patients covered by commercial health insurance may pay **as little as \$0*** copay per fill of DUPIXENT (maximum benefit of \$13,000 per patient per calendar year).

*Approval is not guaranteed. Program has an annual maximum of \$13,000. **THIS IS NOT INSURANCE.** Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed or restricted. **DUPIXENT MyWay** reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.

YOU MAY BE ELIGIBLE FOR THE COPAY CARD PROGRAM IF YOU:

- Have commercial insurance
- Have a DUPIXENT prescription for an FDA-approved condition
- Are a resident of the 50 United States, the District of Columbia, Puerto Rico, Guam, or the US Virgin Islands; and are a patient or caregiver aged 18 years or older

YOU ARE NOT ELIGIBLE IF YOU:

- Have government insurance—Medicaid, Medicare, Veterans Affairs, Department of Defense, TRICARE, other federal or state programs

Not eligible? **DUPIXENT MyWay** may be able to provide additional financial assistance.

SUPPORT YOU DESERVE



The **DUPIXENT MyWay** Copay Card may help eligible patients cover the out-of-pocket cost of DUPIXENT.

ACCESSING COPAY SAVINGS

Eligible patients can access Copay Card benefits in 2 ways:

- Call 1-800-DUPIXENT to ensure Copay benefits are applied to your script.
- Visit **DUPIXENT.com**. Here you can apply for a card online.

Once approved, provide the Copay Card numbers on your card to the specialty pharmacy when they call you to set up delivery of DUPIXENT. The pharmacy will apply the card to reduce your out-of-pocket costs and will note the card number in your record for future refills.

HAVE YOU ALREADY PAID OUT-OF-POCKET COSTS FOR DUPIXENT?

You may be eligible for reimbursement of your out-of-pocket expenses if you meet program eligibility requirements and:

- You were enrolled in the Copay Card Program but did not use it when you filled your DUPIXENT prescription
- You paid a copay for DUPIXENT before enrolling into the Copay Card Program
- Your health plan did not accept your Copay Card

TO APPLY FOR REIMBURSEMENT:

- Visit www.patientrebateonline.com to submit for reimbursement of out-of-pocket costs or to check the card balance. You will be asked to register and login
- Be sure to have ready:
 - Your prescription insurance card
 - Prescription label
 - Your prescription receipt

Please allow 7-10 days to process your request.

Go to DUPIXENT.com/asthma for full program information.

DUPIXENT[®] 
(dupilumab) Injection
200mg · 300mg

CHILDREN & ADULTS CAN DO MORE WITH LESS ASTHMA

DUPIXENT may help you or your child breathe better.

For more information on what
DUPIXENT can do for you, click [here](#).

IMPORTANT SAFETY INFORMATION: Do not use if you are
allergic to dupilumab or to any of the ingredients in DUPIXENT[®].

Please see additional Important Safety Information throughout
and accompanying full [Prescribing Information](#) including
[Patient Information](#).

Reference:

1. Data on file, Sanofi and Regeneron, US. LH Patient View; data through
June 2023.

sanofi | **REGENERON**[®]

Health information contained herein is provided for
general educational purposes only. Your doctor is the
single best source of information regarding your health.
Please consult your doctor if you have any questions
about your health or treatment.

© 2023 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. DUPIXENT[®] and DUPIXENT MyWay[®]
are registered trademarks of Sanofi Biotechnology.
DUP.23.08.0477 08/23