

TALKING TO YOUR TEEN'S ECZEMA SPECIALIST ABOUT DUPIXENT

Growing up with moderate-to-severe eczema (atopic dermatitis) is hard, especially for teens (12-17 years). The unpredictability of flare-ups can be a struggle. And while caring for a teen with eczema has its own set of difficulties, DUPIXENT can help manage their eczema flare-ups even when their skin looks clear. Clearer skin. Less itch. That's a difference your teen can feel inside and out.

Use the information below to help you start a conversation with your doctor about whether DUPIXENT is right for your teen.

WHAT TO SHARE

To get the best care for your teen, it's important to be open about how eczema affects them.

How severe is your teen's eczema? (Scale: 1=not at all severe, 10=extremely severe)

① ② ③ ④ ⑤ ⑥ ⑦ ⑧ ⑨ ⑩

How often does your teen have a flare-up?

Less than once a month 1-2 times a month More than twice a month

How long do your teen's flare-ups typically last?

1-5 days 5-10 days 10-15 days 15+ days

How does your teen's eczema affect them?

WHAT TO ASK

About DUPIXENT

- Could DUPIXENT help my teen?
- What results have you seen with teens on DUPIXENT?
- What are the side effects of DUPIXENT?

Ask your teen's eczema specialist if DUPIXENT is right for them.

Dosing

- How is DUPIXENT taken?
- How often would my teen need to take DUPIXENT?

Support

- Are there programs to help cover the cost of DUPIXENT?
- How have other patients managed the injection process?
- How do I fill a prescription for DUPIXENT?

INDICATION

DUPIXENT is a prescription medicine used to treat people 12 years and older with moderate-to-severe atopic dermatitis (eczema) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids. It is not known if DUPIXENT is safe and effective in children with atopic dermatitis under 12 years of age.

IMPORTANT SAFETY INFORMATION

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT.

**Please see additional Important Safety Information for DUPIXENT on next page.
Please see accompanying full [Prescribing Information](#) including [Patient Information](#).**

IMPORTANT SAFETY INFORMATION (CONT'D)

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have eye problems
- have a parasitic (helminth) infection
- are taking oral, topical, or inhaled corticosteroid medicines. **Do not** stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.
- are scheduled to receive any vaccinations. You should not receive a "live vaccine" if you are treated with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. If you are taking asthma medicines, do not change or stop your asthma medicine without talking to your healthcare provider.

DUPIXENT can cause serious side effects, including:

- **Allergic reactions (hypersensitivity), including a severe reaction known as anaphylaxis.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following symptoms: breathing problems, fever, general ill feeling, swollen lymph nodes, swelling of the face, mouth and tongue, hives, itching, fainting, dizziness, feeling lightheaded (low blood pressure), joint pain, or skin rash.
- **Eye problems.** Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision.

The most common side effects include injection site reactions, eye and eyelid inflammation, including redness, swelling and itching, pain in the throat (oropharyngeal pain) and cold sores in your mouth or on your lips.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed. DUPIXENT is given as an injection under the skin (subcutaneous injection). If your healthcare provider decides that you or a caregiver can give DUPIXENT injections, you or your caregiver should receive training on the right way to prepare and inject DUPIXENT. **Do not** try to inject DUPIXENT until you have been shown the right way by your healthcare provider. In adolescents 12 years of age and older, it is recommended that DUPIXENT be administered by or under supervision of an adult.

Please see accompanying full [Prescribing Information](#) including [Patient Information](#).

Patient Information
DUPIXENT® (DU-pix'-ent)
(dupilumab)
injection, for subcutaneous use

What is DUPIXENT?

DUPIXENT is a prescription medicine used:

- to treat people aged 12 years and older with moderate-to-severe atopic dermatitis (eczema) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids.
- with other asthma medicines for the **maintenance treatment of moderate-to-severe asthma** in people aged 12 years and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing.
- DUPIXENT works by blocking two proteins that contribute to a type of inflammation that plays a major role in atopic dermatitis and asthma.
- DUPIXENT is not used to treat sudden breathing problems
- It is not known if DUPIXENT is safe and effective in children with atopic dermatitis under 12 years of age.
- It is not known if DUPIXENT is safe and effective in children with asthma under 12 years of age.

Do not use DUPIXENT if you are allergic to dupilumab or to any of the ingredients in DUPIXENT. See the end of this leaflet for a complete list of ingredients in DUPIXENT.

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have eye problems (if you also have atopic dermatitis).
- have a parasitic (helminth) infection.
- are taking oral, topical, or inhaled corticosteroid medicines. **Do not** stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.
- are scheduled to receive any vaccinations. You should not receive a "live vaccine" if you are treated with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.

Pregnancy Registry. There is a pregnancy registry for women who take DUPIXENT during pregnancy. The purpose of this registry is to collect information about your health and your baby's health. You can talk to your healthcare provider or contact 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupilixent/> to enroll in this registry or get more information.

- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. If you have asthma and are taking asthma medicines, do not change or stop your asthma medicine without talking to your healthcare provider.

How should I use DUPIXENT?

- **See the detailed "Instructions for Use" that comes with DUPIXENT for information on how to prepare and inject DUPIXENT and how to properly store and throw away (dispose of) used DUPIXENT pre-filled syringes.**
- Use DUPIXENT exactly as prescribed by your healthcare provider.
- DUPIXENT comes as a single-dose pre-filled syringe with needle shield.
- DUPIXENT is given as an injection under the skin (subcutaneous injection).
- If your healthcare provider decides that you or a caregiver can give the injections of DUPIXENT, you or your caregiver should receive training on the right way to prepare and inject DUPIXENT. **Do not** try to inject DUPIXENT until you have been shown the right way by your healthcare provider. In adolescents 12 years of age and older, it is recommended that DUPIXENT be administered by or under supervision of an adult.
- If you miss a dose of DUPIXENT, give the injection within 7 days from the missed dose, then continue with the original schedule. If the missed dose is not given within 7 days, wait until the next scheduled dose to give your DUPIXENT injection.
- If you inject more DUPIXENT than prescribed, call your healthcare provider right away.
- Your healthcare provider may prescribe other medicines to use with DUPIXENT. Use the other prescribed medicines exactly as your healthcare provider tells you to.

What are the possible side effects of DUPIXENT?

DUPIXENT can cause serious side effects, including:

- **Allergic reactions (hypersensitivity), including a severe reaction known as anaphylaxis.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following symptoms:

- | | | |
|-----------------------|---|---|
| ○ breathing problems | ○ swelling of the face, mouth, and tongue | ○ fainting, dizziness, feeling lightheaded (low blood pressure) |
| ○ fever | ○ hives | ○ joint pain |
| ○ general ill feeling | ○ itching | ○ skin rash |
| ○ swollen lymph nodes | | |

- **Eye problems.** If you have atopic dermatitis, tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision.

- **Inflammation of your blood vessels.** Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. It is not known whether this is caused by DUPIXENT. Tell your healthcare provider right away if you have:

- | | |
|-----------------------|--|
| ○ rash | ○ chest pain |
| ○ shortness of breath | ○ a feeling of pins and needles or numbness of your arms or legs |
| ○ persistent fever | |

The most common side effects of DUPIXENT include:

- injection site reactions
- eye and eyelid inflammation, including redness, swelling, and itching (if you also have atopic dermatitis)
- pain in the throat (oropharyngeal pain)
- cold sores in your mouth or on your lips

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of DUPIXENT.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of DUPIXENT.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use DUPIXENT for a condition for which it was not prescribed. Do not give DUPIXENT to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about DUPIXENT that is written for health professionals.

What are the ingredients in DUPIXENT?

Active ingredient: dupilumab

Inactive ingredients: L-arginine hydrochloride, L-histidine, polysorbate 80, sodium acetate, sucrose, and water for injection.

REGENERON SANOFI GENZYME 

Manufactured by: Regeneron Pharmaceuticals, Inc., Tarrytown, NY 10591 U.S. License No. 1760

Marketed by: sanofi-aventis U.S. LLC (Bridgewater, NJ 08807) and Regeneron Pharmaceuticals, Inc. (Tarrytown, NY 10591)

DUPIXENT® is a registered trademark of Sanofi Biotechnology / © 2019 Regeneron Pharmaceuticals, Inc. / sanofi-aventis U.S. LLC. All rights reserved.

For more information about DUPIXENT, go to www.DUPIXENT.com or call 1- 844-DUPIXENT (1-844-387-4936).

This Patient Information has been approved by the U.S. Food and Drug Administration.

Issued: March 2019

US-DUP-1126(1)