

Living as an adult with uncontrolled chronic rhinosinusitis with nasal polyps (CRSwNP) can be challenging. That's why it's important to get the conversation started with your doctor so you can discuss appropriate treatment options.

SHARE

your thoughts about

- Treatments you've tried and are currently using
- Inhaled/Oral steroid use
- Your willingness to try a new treatment
- History of surgery

DISCUSS

the symptoms you've experienced and how they've affected you.

- Nasal congestion
- Difficulty breathing through your nose
- Loss of taste/smell
- Chronic sinus pressure/pain

These symptoms affect me when/because

ASK

about DUPIXENT and if it's right for you.

- How is DUPIXENT different from other treatment options?
- How does DUPIXENT work?
- How would I take DUPIXENT?
- How do I get started with DUPIXENT?
- What are the possible side effects of DUPIXENT?
- How can DUPIXENT help treat my uncontrolled CRSwNP?

INDICATION

DUPIXENT is a prescription medicine used with other medicines to treat chronic rhinosinusitis with nasal polyposis in adults whose disease is not controlled. It is not known if DUPIXENT is safe and effective in children with chronic rhinosinusitis with nasal polyposis under 18 years of age.

IMPORTANT SAFETY INFORMATION

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT.

Please see additional Important Safety Information for DUPIXENT on next page.
Please see accompanying full [Prescribing Information](#) and [Patient Information](#).

IMPORTANT SAFETY INFORMATION (cont'd)

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have eye problems.
- have a parasitic (helminth) infection.
- are taking oral, topical, or inhaled corticosteroid medicines. Do not stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. If you are taking asthma medicines, do not change or stop your asthma medicine without talking to your healthcare provider.

DUPIXENT can cause serious side effects, including:

- **Allergic reactions (hypersensitivity), including a severe reaction known as anaphylaxis.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following symptoms: breathing problems, fever, general ill feeling, swollen lymph nodes, swelling of the face, mouth and tongue, hives, itching, fainting, dizziness, feeling lightheaded (low blood pressure), joint pain, or skin rash.
- **Eye problems.** Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision.
- **Inflammation of your blood vessels.** Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. It is not known whether this is caused by DUPIXENT. Tell your healthcare provider right away if you have: rash, shortness of breath, persistent fever, chest pain, or a feeling of pins and needles or numbness of your arms or legs.

The most common side effects in patients with chronic rhinosinusitis with nasal polyposis include injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, high count of a certain white blood cell (eosinophilia), trouble sleeping (insomnia), toothache, gastritis and joint pain (arthralgia).

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed. DUPIXENT is given as an injection under the skin (subcutaneous injection). If your healthcare provider decides that you or a caregiver can give DUPIXENT injections, you or your caregiver should receive training on the right way to prepare and inject DUPIXENT. **Do not** try to inject DUPIXENT until you have been shown the right way by your healthcare provider.

Please see accompanying full [Prescribing Information](#) including [Patient Information](#).

Patient Information
DUPIXENT® (DU-pix'-ent)
(dupilumab)
injection, for subcutaneous use

What is DUPIXENT?

DUPIXENT is a prescription medicine used:

- to treat people aged 12 years and older with moderate-to-severe atopic dermatitis (eczema) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids.
- with other asthma medicines for the maintenance treatment of moderate-to-severe asthma in people aged 12 years and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing.
- with other medicines to treat chronic rhinosinusitis with nasal polyposis in adults whose disease is not controlled.
- DUPIXENT works by blocking two proteins that contribute to a type of inflammation that plays a major role in atopic dermatitis, asthma, and chronic rhinosinusitis with nasal polyposis.
- DUPIXENT is not used to treat sudden breathing problems.
- It is not known if DUPIXENT is safe and effective in children with atopic dermatitis or asthma under 12 years of age.
- It is not known if DUPIXENT is safe and effective in children with chronic rhinosinusitis with nasal polyposis under 18 years of age.

Do not use DUPIXENT if you are allergic to dupilumab or to any of the ingredients in DUPIXENT. See the end of this leaflet for a complete list of ingredients in DUPIXENT.

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have eye problems.
- have a parasitic (helminth) infection.
- are taking oral, topical, or inhaled corticosteroid medicines. **Do not** stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” if you are treated with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.

Pregnancy Registry. There is a pregnancy registry for women who take DUPIXENT during pregnancy. The purpose of this registry is to collect information about your health and your baby’s health. You can talk to your healthcare provider or contact 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/> to enroll in this registry or get more information.

- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.
- Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. If you have asthma and are taking asthma medicines, do not change or stop your asthma medicine without talking to your healthcare provider.

How should I use DUPIXENT?

- **See the detailed “Instructions for Use” that comes with DUPIXENT for information on how to prepare and inject DUPIXENT and how to properly store and throw away (dispose of) used DUPIXENT pre-filled syringes.**
- Use DUPIXENT exactly as prescribed by your healthcare provider.
- DUPIXENT comes as a single-dose pre-filled syringe with needle shield.
- DUPIXENT is given as an injection under the skin (subcutaneous injection).
- If your healthcare provider decides that you or a caregiver can give the injections of DUPIXENT, you or your caregiver should receive training on the right way to prepare and inject DUPIXENT. **Do not** try to inject DUPIXENT until you have been shown the right way by your healthcare provider. In children 12 years of age and older, it is recommended that DUPIXENT be given by or under the supervision of an adult.
- If you miss a dose of DUPIXENT, give the injection within 7 days from the missed dose, then continue with the original schedule. If the missed dose is not given within 7 days, wait until the next scheduled dose to give your DUPIXENT injection.
- If you inject more DUPIXENT than prescribed, call your healthcare provider right away.
- Your healthcare provider may prescribe other medicines to use with DUPIXENT. Use the other prescribed medicines exactly as your healthcare provider tells you to.

What are the possible side effects of DUPIXENT?

DUPIXENT can cause serious side effects, including:

- **Allergic reactions (hypersensitivity), including a severe reaction known as anaphylaxis.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following symptoms:
 - breathing problems
 - swelling of the face, mouth, and tongue
 - fainting, dizziness, feeling lightheaded (low blood pressure)
 - fever
 - hives
 - joint pain
 - general ill feeling
 - itching
 - skin rash
 - swollen lymph nodes
- **Eye problems.** Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision.

• **Inflammation of your blood vessels.** Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. It is not known whether this is caused by DUPIXENT. Tell your healthcare provider right away if you have:

- rash
- shortness of breath
- persistent fever
- chest pain
- a feeling of pins and needles or numbness of your arms or legs

The most common side effects of DUPIXENT include:

- injection site reactions
- eye and eyelid inflammation, including redness, swelling, and itching
- pain in the throat (oropharyngeal pain)
- cold sores in your mouth or on your lips
- high count of a certain white blood cell (eosinophilia)
- trouble sleeping (insomnia)
- toothache
- gastritis
- joint pain (arthralgia)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of DUPIXENT.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of DUPIXENT.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use DUPIXENT for a condition for which it was not prescribed. Do not give DUPIXENT to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about DUPIXENT that is written for health professionals.

What are the ingredients in DUPIXENT?

Active ingredient: dupilumab

Inactive ingredients: L-arginine hydrochloride, L-histidine, polysorbate 80, sodium acetate, sucrose, and water for injection.

REGENERON **SANOBI GENZYME** 

Manufactured by: Regeneron Pharmaceuticals, Inc., Tarrytown, NY 10591 U.S. License No. 1760

Marketed by: sanofi-aventis U.S. LLC (Bridgewater, NJ 08807) and Regeneron Pharmaceuticals, Inc. (Tarrytown, NY 10591)

DUPIXENT® is a registered trademark of Sanofi Biotechnology / © 2019 Regeneron Pharmaceuticals, Inc. / sanofi-aventis U.S. LLC. All rights reserved.

For more information about DUPIXENT, go to www.DUPIXENT.com or call 1-844-DUPIXENT (1-844-387-4936).

This Patient Information has been approved by the U.S. Food and Drug Administration.

Issued: June 2019