

DUPIXENT[®]
(dupilumab) Injection
200mg · 300mg



**EoE SYMPTOMS
CAUSING TROUBLE WITH FOOD?
START TREATING WITH
DUPIXENT**

Patient Portrayal

The only FDA-approved treatment of its kind for
EoE patients ages 1 and up who weigh at least 33 lbs

INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 1 year of age and older with eosinophilic esophagitis (EoE), who weigh at least 33 pounds (15 kg). It is not known if DUPIXENT is safe and effective in children with EoE under 1 year of age, or who weigh less than 33 pounds (15 kg).

IMPORTANT SAFETY INFORMATION

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT[®].

Please see additional **Important Safety Information** throughout and full **Prescribing Information** including **Patient Information**.

EoE = eosinophilic esophagitis.



WHAT IS EoE?

EoE, or eosinophilic esophagitis, is a **chronic and progressive condition** that can affect both children and adults, where inflammation in the esophagus causes it to narrow, often making it very difficult to swallow.

The inflammation occurs when a large number of immune cells (including eosinophils) build up within the esophagus. This buildup is due to an abnormal response of the immune system, known as Type 2 inflammation.



Normal esophagus



Immune cell buildup causes inflammation



Inflammation can lead to difficulty swallowing, scarring, and other complications from EoE

EoE is a long-term and progressive condition, it's important to find a treatment path for children early. The longer it goes untreated, the more damage it can do.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a "live vaccine" right before and during treatment with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
 - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.



Patient Portrayal

EoE CAN BE HARD TO DIAGNOSE

EoE symptoms can look like other conditions such as food allergies or GERD (gastroesophageal reflux disease). This makes diagnosis challenging and can lead to treatment delays.

To confirm EoE, a gastroenterologist will perform tests such as an endoscopy with biopsy. This helps identify if there are increased levels of eosinophils in the esophagus.

If it's EoE, you and your child's doctor will customize a treatment plan based on your child's needs.



REMEMBER:

Following your child's treatment plan consistently is key to managing EoE and helping your child feel better.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#) including [Patient Information](#).



LIVING WITH EoE

CAREGIVERS AND CHILDREN OFTEN ADAPT THEIR MEALTIME BEHAVIORS TO HELP MANAGE EoE



When your child has EoE symptoms, you both find ways to get through meals more easily. Your child may have trouble or avoid certain foods. You may notice coping behaviors such as **excessive chewing, being uncooperative around food, and slow eating.**

So as their caregiver, you adapt. Maybe it's cutting their food in tiny pieces or covering it in sauce to make it go down easier. This can be stressful and a lot of work for you, but you're doing what you can to make mealtimes a smoother experience.

Now you're taking an important step by exploring options that can actually help treat EoE, not just dealing with the symptoms.

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

WHAT ARE THE SIGNS AND SYMPTOMS OF EoE?

THESE ARE SOME OF THE MOST COMMON ONES TO LOOK FOR:

- Food refusal
- Vomiting
- Regurgitation
- Heartburn (burning feeling in chest)
- Acid reflux (acid coming up from stomach into throat)
- Difficulty swallowing
- Food getting stuck in the esophagus (impaction)

RECOGNIZE ANY?

It's important to bring these up with your child's Gastroenterologist or Allergist.

“My younger daughter would just choke and throw up all the time. When she was a toddler, she would throw up everything.”

–Sara F., mother of Liv, real DUPIXENT EoE patient
Sponsored partner



[Click here for caregiver tools & tips.](#)



Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.



DUPIXENT 
(dupilumab) Injection
200mg • 300mg


DUPIXENT[®] (dupilumab) PROACTIVELY TREATS EoE FOR CHILDREN

AS YOUNG AS 1 YEAR OLD (≥33 LBS)

DUPIXENT is a biologic that **helps block two of the key sources of inflammation** inside the body that can cause EoE. This inflammation is driven by immune cells, including eosinophils.

WHAT ARE EOSINOPHILS?

Eosinophils are part of your immune system, but when there are too many in the esophagus, it results in inflammation that can trigger conditions like EoE. Your child's doctor collects these cells through a biopsy to measure your child's inflammation.



A UNIQUE TREATMENT APPROACH



DUPIXENT:

- TARGETS AN UNDERLYING SOURCE OF TYPE 2 INFLAMMATION
- IS NOT A STEROID
- IS NOT AN IMMUNOSUPPRESSANT

“Within weeks I noticed that Noah’s symptoms were less severe. He wasn’t complaining of stomach pain or getting food stuck as often.”

–Britt B., mother of Noah,
real DUPIXENT EoE patient
Individual results may vary
Sponsored partner



[Click here to learn more about Britt’s experience with her son’s EoE and how DUPIXENT helped.](#)

IMPORTANT SAFETY INFORMATION (CONT'D)

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have EoE and asthma and use an asthma medicine. **Do not** change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

Please see additional [Important Safety Information](#) throughout and full [Prescribing Information](#) including [Patient Information](#).



DUPIXENT
(dupilumab) Injection
200mg · 300mg

WHY YOU SHOULD GO WITH DUPIXENT[®] (dupilumab)



Patient Portrayal


IMPORTANT SAFETY INFORMATION (CONT'D)

DUPIXENT can cause serious side effects, including:

- Allergic reactions. DUPIXENT can cause allergic reactions, including skin reactions, that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye or painful red or blue bumps under the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

DUPIXENT IS PROVEN TO REDUCE INFLAMMATION IN THE ESOPHAGUS*

Significant percentage of patients had their eosinophils reduced to 6 or less.*



In an analysis conducted after the trial, patients had a decrease in their average esophageal eosinophil count while patients on placebo had an increase.*

84% DECREASE ON DUPIXENT **20% INCREASE ON PLACEBO**

Analysis was performed on patients on FDA-approved dosing of DUPIXENT in patients aged 1-11 years, weighing at least 33 lbs.

Analysis was not evaluated during the clinical trial, and any results may be due to chance. Individual results may vary.

Visible improvements of the esophagus were seen in children treated with DUPIXENT.*



*Results at 4 months from 61 patients ages 1 to 11 years old (≥33 lbs) with EoE in a two-part study.

“Some time after Paisley started DUPIXENT, we started to see improvements in her EoE symptoms. Gradually, she choked less and less.”

–Ali P., mother of Paisley, real DUPIXENT EoE patient

Individual results may vary
Sponsored partner



Please see additional Important Safety Information throughout and full [Prescribing Information](#) including [Patient Information](#).



DUPIXENT[®]
(dupilumab) Injection
200mg • 300mg

CAREGIVERS SAW DUPIXENT[®] (dupilumab) HELPS RELIEVE EoE SYMPTOMS THAT GIVE KIDS TROUBLE WITH FOOD*†

*Results at 4 months from 61 patients ages 1 to 11 years old (≥33 lbs) with EoE in a two-part study.

†As measured by days with symptoms using the Pediatric EoE Signs/Symptoms Questionnaire-Caregiver (PESQ-C).



Patient Portrayal

IMPORTANT SAFETY INFORMATION (CONT'D)

DUPIXENT can cause serious side effects, including:

- **Joint aches and pain.** Some people who use **DUPIXENT** have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop **DUPIXENT** if you develop joint symptoms.

DUPIXENT TARGETS AN UNDERLYING SOURCE OF INFLAMMATION THAT CONTRIBUTES TO EoE



DUPIXENT is the first and only FDA-approved treatment of its kind for EoE in children as young as 1 year old (≥33 lbs).

There are other treatments for EoE. These include:

- Food elimination diets
- Swallowed corticosteroids
- Proton pump inhibitors (PPIs)

Some of these treatments may provide short-term relief. Yet for many patients, their EoE symptoms are still uncontrolled.

DUPIXENT is neither a PPI nor a swallowed topical corticosteroid. PPIs aren't FDA-approved to treat EoE.



“A few months after we started **DUPIXENT**, we realized that our son's EoE symptoms, like regurgitating food during meals, had improved.”

–Jeni B., mother of Josh, real **DUPIXENT** EoE patient

Individual results may vary
Sponsored partner



IMPORTANT SAFETY INFORMATION (CONT'D)

The most common side effects in patients with EoE include injection site reactions, upper respiratory tract infections, cold sores in your mouth or on your lips, and joint pain (arthralgia).

Please see additional **Important Safety Information** throughout and full **Prescribing Information** including **Patient Information**.



ASK THEIR DOCTOR

These questions will help you prepare for your child's upcoming appointment. Complete the questionnaire with your child, then bring it with you to discuss with their Gastroenterologist or Allergist.

I first noticed my child's symptoms of eosinophilic esophagitis (EoE) at _____ months/years old.

SYMPTOMS

- Bellyache
- Heartburn (burning feeling in chest)
- Acid reflux (acid coming up from stomach into throat)
- Regurgitation (food coming up from stomach into mouth)
- Vomiting (throwing up)
- Food refusal (refusing to eat a meal)
- Trouble swallowing food
- Food stuck in throat

THEY EXPERIENCE SYMPTOMS:

- A few times a year Monthly Weekly
- Daily Other (e.g., "twice a week"): _____

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and full [Prescribing Information](#) including [Patient Information](#).

HOW OFTEN DOES YOUR CHILD AVOID FOOD BECAUSE THEY'RE AFRAID OF EXPERIENCING EoE SYMPTOMS?

- A few times a year Monthly Weekly
- Daily Other (e.g., "twice a week"): _____

WHICH OF THESE STRATEGIES HAVE YOU TRIED FOR YOUR CHILD IN THE PAST YEAR?

- Modifying foods (e.g., putting ingredients in the blender or cutting up food into very small pieces)
- Drinking a lot of liquid
- Adding sauces to foods
- Eating slowly
- A proton pump inhibitor (PPI) as prescribed
- Avoiding certain foods/changing my child's diet
- A swallowed topical corticosteroid (TCS) as prescribed
- Other: _____



THINGS TO THINK ABOUT

My child's eosinophilic esophagitis (EoE) affects our family by _____

My goals for the management of my child's EoE are _____

How soon would my child start to feel the effects of DUPIXENT?

[Click here for tips on having an even more productive conversation with your child's doctor.](#)



HOW DUPIXENT[®] (dupilumab) IS TAKEN

DUPIXENT is an injection given subcutaneously (under the skin). If your child's healthcare provider decides you can inject your child with DUPIXENT, they will train you on the right way to prepare and inject it, at a frequency based on the chart below.

In children 1 to <12 years of age, DUPIXENT should be given by a caregiver. In children 12+ years, it's recommended DUPIXENT be administered by or under supervision of an adult.



PRE-FILLED PEN

With a Pre-filled Pen, your child won't have to see the needle. A viewing window and two click sounds let you see and hear when the dose is being delivered.

Available in 200 mg and 300 mg.

PRE-FILLED SYRINGE

With the Pre-filled Syringe, you control the injection speed. The contoured finger grip provides a firm hold, and the device features a needle shield to protect against accidental sticks.

Available in 200 mg and 300 mg.

WEIGHT

33 to <66 lbs

66 to <88 lbs

88 lbs+

FREQUENCY

200 mg every 2 weeks

300 mg every 2 weeks

300 mg every week

IMPORTANT SAFETY INFORMATION (CONT'D)

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 1 to less than 12 years of age, DUPIXENT should be given by a caregiver.

WHAT CAN DUPIXENT MyWay[®] DO FOR YOU?

PAY AS LITTLE AS \$0*

The **DUPIXENT MyWay Copay Card** may help reduce the out-of-pocket cost of DUPIXENT. If you have commercial health insurance, you may be eligible to pay as little as \$0* in copay per fill of DUPIXENT through the **DUPIXENT MyWay Copay Card**.

[Click here to see if you're eligible.](#)



SUPPORT FOR EVERY STEP

Once your child has been prescribed DUPIXENT, the **DUPIXENT MyWay patient support program is here to help**. It offers all kinds of additional support, like supplemental injection training from a Nurse Educator, refill and injection reminders, and tips and tools to help manage treatment.

*Subject to the program maximum per patient per calendar year. Approval is not guaranteed. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs, including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed, or restricted. **DUPIXENT MyWay** reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT.

Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.

Please see additional **Important Safety Information throughout and full Prescribing Information including Patient Information.**

DUPIXENT[®]
(dupilumab) Injection
200mg · 300mg



**WITH YOUR CHILD'S EoE
UNDER CONTROL, THINK
OF THE POSSIBILITIES**

Ask your child's EoE specialist
about DUPIXENT.

Patient Portrayal

IMPORTANT SAFETY INFORMATION

Do not use if you are allergic to dupilumab
or to any of the ingredients in DUPIXENT.

**Please see additional Important Safety
Information throughout and full Prescribing
Information including Patient Information.**

sanofi | **REGENERON**[®]

© 2026 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. DUPIXENT and DUPIXENT MyWay are registered
trademarks of Sanofi or an affiliate. US.DUP.25.09.0237 05/26

Health information contained herein is provided for general educational
purposes only. Your doctor is the single best source of information
regarding your health. Please consult your doctor if you have any
questions about your health or treatment.