



DU MORE WITH LESS ASTHMA

APPROVED FOR AGES 6 AND UP

#1 PRESCRIBED BIOLOGIC FOR ASTHMA¹

INDICATION: DUPIXENT is a prescription medicine used with other asthma medicines for the maintenance treatment of moderate-to-severe eosinophilic or oral steroid dependent asthma in adults and children 6 years of age and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing. DUPIXENT is not used to relieve sudden breathing problems and will not replace an inhaled rescue medicine. It is not known if DUPIXENT is safe and effective in children with asthma under 6 years of age.

Please see additional Important Safety Information throughout and full [Prescribing Information](#) including [Patient Information](#).

DUPIXENT 
(dupilumab) Injection
200mg • 300mg

KNOW MORE ABOUT ASTHMA

ASTHMA IS A RESPIRATORY DISEASE,
USUALLY CHARACTERIZED BY CHRONIC AIRWAY
INFLAMMATION. SYMPTOMS MAY VARY OVER
TIME AND CAN BE REVERSIBLE.

SYMPTOMS CAN INCLUDE:



SHORTNESS
OF BREATH



CHEST
TIGHTNESS



WHEEZING



COUGHING


WHAT CAN CAUSE ASTHMA SYMPTOMS?

Some of the most common asthma triggers are:

- exercise
- respiratory infections
- pet dander or dust
- irritants like cleaning products or perfume
- weather conditions

What's more, **underlying inflammation can still be there even when you don't have symptoms.**

With treatment, this inflammation in the lungs may be reduced, allowing you to breathe better.



IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you: have a parasitic (helminth) infection; are scheduled to receive any vaccinations. You should not receive a "live vaccine" right before and during treatment with DUPIXENT; are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby. A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>; are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see additional Important Safety Information throughout and full [Prescribing Information](#) including [Patient Information](#).

BREATHING MADE BETTER

WHAT DUPIXENT CAN DO:



**BLOCK A KEY SOURCE OF INFLAMMATION
IN THE LUNGS**



HELP PREVENT ASTHMA ATTACKS



**HELP PEOPLE WITH ASTHMA ACHIEVE
LONG-LASTING BREATHING IMPROVEMENT**



REDUCE OR ELIMINATE ORAL STEROID USE

And DUPIXENT requires only **1-2 doses a month***

81% EXPERIENCED
FEWER
ASTHMA ATTACKS^{†‡}

*Dosing based on weight and age.

†In those with high eosinophils (a type of white blood cell).

‡The annual rate of severe asthma attacks was 0.20 with DUPIXENT 300 mg vs 1.04 with placebo.

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

HOW IT WORKS

BEFORE DUPIXENT



The chronic underlying inflammation in asthma may cause airways to become constricted and narrow, making it harder to breathe.

AFTER DUPIXENT



DUPIXENT helps reduce underlying inflammation.

Illustrative image of DUPIXENT mechanism only, and not reflective of treatment effect.

HOW IS DUPIXENT DIFFERENT?


- It's a prescription biologic that targets an underlying cause of asthma in the lungs
- It's not a steroid or inhaler

IMPORTANT SAFETY INFORMATION: Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or use an asthma medicine. **Do not** change or stop your other medicines, including corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.


REAL PEOPLE, REAL RESULTS

**PRESCRIBED TO OVER 150,000 ASTHMA PATIENTS
AND COUNTING!**^{1*†}



Pamela

Real DUXENT patient



Lydia

Real DUXENT patient

Individual results may vary.

*LH Patient View: data through September 2024.

†New adult and pediatric patients aged 6 years and older with moderate-to-severe eosinophilic or OCS-dependent asthma. OCS=oral corticosteroids

DU MORE YOU

DUXENT targets a key source of chronic airway inflammation that makes asthma difficult to control.

In multiple clinical studies, people taking DUXENT saw improvements, including:

- LESS SHORTNESS OF BREATH
- LESS WHEEZING
- NOT WAKING AT NIGHT
- INCREASED ABILITY TO DO NORMAL ACTIVITIES

The relationship between DUXENT and these results was studied but not definitively proven.

The most common side effects include injection site reactions, pain in the throat (oropharyngeal pain), and high count of a certain white blood cell (eosinophilia), with the addition of parasitic (helminth) infections observed in children.

IMPORTANT SAFETY INFORMATION: DUXENT can cause serious side effects, including: Allergic reactions. DUXENT can cause allergic reactions that can sometimes be severe.

Stop using DUXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.

Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.



KIDS CAN DU MORE, TOO

Let them play! DUPIXENT was proven to help improve lung function and prevent asthma attacks in children 6+ years of age.



DUPIXENT can help your child breathe better in **as little as 2 weeks** and achieve **long-lasting breathing improvement**



Children who used DUPIXENT to treat their asthma **reduced their use of oral steroids**



DUPIXENT is just **1 or 2 doses a month***

*Dosing based on weight and age.

We're happy with the decision we made. It's worth Aston taking an injection to help him breathe better and do more of the things he loves.



Aston, age 9

Christan

Real DUPIXENT patient caregiver/
Aston's mom

IMPORTANT SAFETY INFORMATION: Inflammation of your blood vessels. Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. Tell your healthcare provider right away if you have: rash, chest pain, worsening shortness of breath, brown or dark colored urine, persistent fever, or a feeling of pins and needles or numbness of your arms or legs.

Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.

RELYING ON ORAL STEROIDS?

While they may be helpful when prescribed, long-term use of oral steroids can cause:

- WEIGHT GAIN
- MOOD CHANGE
- HIGH BLOOD PRESSURE
- DIABETES
- VISION PROBLEMS

This may not be an exhaustive list of side effects associated with oral steroid use.

I had to make sure my bag was packed with my daily steroids that I needed to get through the day. Since I started DUPIXENT, I've been able to decrease my reliance on oral steroids.



Tim
Real DUPIXENT patient

IN A STUDY OF PEOPLE DEPENDENT ON STEROIDS:

86% OF PEOPLE ON DUPIXENT REDUCED OR ELIMINATED THEIR STEROID USE*†

*In a clinical study of patients 12+ years old with DUPIXENT 300 mg + standard of care (SOC) vs **68% with placebo + SOC**.

†Do not stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.

READY TO TRY SOMETHING ELSE?

USING THE RULE OF 2
CAN HELP YOU DETERMINE HOW WELL YOUR ASTHMA IS CONTROLLED



- YOU USE A RESCUE INHALER MORE THAN **2X** A WEEK
- YOU HAD **2 OR MORE** COURSES OF ORAL STEROIDS IN THE LAST YEAR
- YOU HAD **2 OR MORE** ASTHMA ATTACKS IN THE LAST YEAR
- YOU WAKE UP AT NIGHT **2 OR MORE** TIMES A MONTH DUE TO ASTHMA

Talk to your doctor about your asthma and ask if DUPIXENT could be right for you.

IMPORTANT SAFETY INFORMATION: Psoriasis. This can happen in people with asthma who receive DUPIXENT. Tell your healthcare provider about any new skin symptoms. Your healthcare provider may send you to a dermatologist for an examination if needed.

Joint aches and pain. Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.



IMPORTANT SAFETY INFORMATION:

The most common side effects in patients with asthma include injection site reactions, high count of a certain white blood cell (eosinophilia), pain in the throat (oropharyngeal pain), and parasitic (helminth) infections.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.

TALK TO YOUR DOCTOR

Use these questions to organize your thoughts and bring the answers to your next appointment.

START WITH THE RULE OF 2 TO DETERMINE YOUR LEVEL OF CONTROL



1. Have you used your rescue inhaler **more than 2X** during a typical week?

Yes

No

2. Did you use **2 or more** courses of oral steroids in the last year?

Yes

No

3. Did you have **2 or more** asthma attacks in the last year?

Yes

No

4. Have you woken up **more than 2X** at night in the past month because of your asthma?

Yes

No

5. How long have you been experiencing asthma symptoms?

_____ Year(s) _____ Month(s)

6. What treatments have you or your loved one been prescribed for asthma?

Select all that apply:

Inhalers

Nebulizers

Biologics

Inhaled steroids

Inhaled long-acting beta antagonists

Oral steroids

None

Other _____

7. How often do these symptoms occur?

Coughing

Never Rarely Occasionally Frequently

Wheezing

Never Rarely Occasionally Frequently

Rapid breathing

Never Rarely Occasionally Frequently

Chest tightness

Never Rarely Occasionally Frequently

8. Have you limited or missed an activity?

Yes No

9. Have you noticed any of the following triggers for your asthma?

Dust mites Air pollution Pets Mold

Perfumes Scented soaps Physical exercise

Other _____

IMPORTANT SAFETY INFORMATION: Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see additional Important Safety Information throughout and full Prescribing Information including Patient Information.

CONNECT WITH AN ASTHMA MENTOR

The **DUPIXENT Mentor Program** is designed to help new patients know what to expect by talking one-on-one with a real DUPIXENT patient mentor.

You can:

-  **LEARN ABOUT THEIR DUPIXENT EXPERIENCE**
-  **ASK QUESTIONS**
-  **GET HELPFUL TIPS**

As long as you've discussed DUPIXENT with your doctor or have been prescribed DUPIXENT, you can register for a mentor so you can hear about their journey and experience with DUPIXENT.



Scan or [click here](#) to request a mentor.


It's easy to lose hope when you have difficulty breathing. I want others to know that I was once there, and wish I had someone to talk to in order to know that there is hope and a chance to breathe better.



Rachel
DUPIXENT MyWay® Patient Mentor

SUPPORT YOU DESERVE

The **DUPIXENT MyWay**[®] Copay Card may help eligible patients cover the out-of-pocket cost of DUPIXENT.



ACCESSING YOUR COPAY CARD

You can access your Copay Card in 2 ways:



Call **1-855-726-4702** to ensure Copay benefits are applied to your script



Visit **DUPIXENT.com** to apply for a card online

Once approved, provide the Copay Card numbers on your card to the specialty pharmacy when they call you to set up delivery of DUPIXENT. The pharmacy will apply the card to reduce your out-of-pocket costs and will note the card number in your record for future refills.

YOUR COPAY MAY BE AS LITTLE AS \$0*

99% OF COMMERCIALLY INSURED PATIENTS
ARE NATIONALLY COVERED FOR DUPIXENT!

Eligible patients with commercial health insurance may pay **as little as \$0*** in copay per fill of DUPIXENT[®] (dupilumab). Terms and conditions apply.

*Subject to the program maximum per patient per calendar year. Approval is not guaranteed. **THIS IS NOT INSURANCE.** Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs, including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed, or restricted. **DUPIXENT MyWay** reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.

YOU MAY BE ELIGIBLE FOR THE COPAY CARD PROGRAM IF YOU:

- Have commercial insurance
- Have a DUPIXENT prescription for an FDA-approved condition
- Are a resident of the 50 United States, the District of Columbia, Puerto Rico, Guam, or the US Virgin Islands, and are a patient or caregiver aged 18 years or older

YOU ARE NOT ELIGIBLE IF YOU:

- Have government insurance—Medicaid, Medicare, Veterans Affairs, Department of Defense, TRICARE, other federal or state programs

NOT ELIGIBLE?

DUPIXENT MyWay[®] may be able to provide additional financial assistance.

HAS YOUR DOCTOR PRESCRIBED DUPIXENT[®] (dupilumab)?

Your healthcare provider is your best resource for information about DUPIXENT and should be your primary point of contact throughout your treatment journey.

YOU BEGIN YOUR JOURNEY.

Your healthcare provider prescribes DUPIXENT, and you enroll into *DUPIXENT MyWay*[®].

- Your healthcare provider can download the enrollment form on **DUPIXENT.com**, help you fill it out, and fax it back to *DUPIXENT MyWay* at **1-844-387-9370**
- You or your healthcare provider can call **1-844-DUPIXEN(T), Option 1**
- Providing your email address allows *DUPIXENT MyWay* to give you more support resources about DUPIXENT

QUESTIONS ABOUT INJECTING? RESOURCES ARE HERE FOR YOU.



Take advantage of our **supplemental injection resources** and **training videos**, scan the QR code or [click here](#).

WHAT CAN *DUPIXENT MyWay*[®] DO FOR YOU?

Enrolling in *DUPIXENT MyWay* can help ensure you receive DUPIXENT as quickly as possible and receive additional support along your treatment journey. For eligible patients, *DUPIXENT MyWay* can:



Help research and explain your insurance benefits



Find ways to help with the out-of-pocket cost of DUPIXENT if you need help with your copay, you are denied coverage from your health plan, or you do not have or experience a loss in insurance



Connect you with a Nurse Educator to answer questions about DUPIXENT and help you start and stay on track with therapy



Connect you with your specialty pharmacy to make sure your prescription is shipped to your home or preferred location



Explain how to properly store DUPIXENT when you receive your shipment



Provide supplemental injection training—in person, virtually, or over the phone—to help you or a caregiver get more familiar with injecting DUPIXENT



Offer a needle-disposing kit or sharps container for proper disposal



Remind you when it is time to refill your DUPIXENT prescription

DUPIXENT 
(dupilumab) Injection
200mg · 300mg



DU MORE WITH LESS ASTHMA

DUPIXENT. Better breathing for you or your child (6+ years).



Scan this QR code or [click here](#)
to get more information on what
DUPIXENT can do for you.

IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Please see additional Important Safety Information throughout and full [Prescribing Information](#) including [Patient Information](#).

Reference:

1. IQVIA NBRx data through March 2025.

sanofi | **REGENERON**®

Health information contained herein is provided for general educational purposes only. Your doctor is the single best source of information regarding your health. Please consult your doctor if you have any questions about your health or treatment.

© 2025 Sanofi and Regeneron Pharmaceuticals, Inc. All Rights Reserved. DUPIXENT® and DUPIXENT MyWay® are registered trademarks of Sanofi Biotechnology. US.DUP.25.04.0478 04/2025