

DU MORE WITH LESS ASTHMA

APPROVED FOR AGES 6 YEARS AND UP



#1 PRESCRIBED BIOLOGIC FOR ASTHMA¹

INDICATION: DUPIXENT is a prescription medicine used with other asthma medicines for the maintenance treatment of moderate-to-severe eosinophilic or oral steroid dependent asthma in adults and children 6 years of age and older whose asthma is not controlled with their current asthma medicines. DUPIXENT helps prevent severe asthma attacks (exacerbations) and can improve your breathing. DUPIXENT may also help reduce the amount of oral corticosteroids you need while preventing severe asthma attacks and improving your breathing. **DUPIXENT is not used** to relieve sudden breathing problems and will not replace an inhaled rescue medicine. It is not known if DUPIXENT is safe and effective in children with asthma under 6 years of age.

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

DUPIXENT® 
(dupilumab) Injection
200mg · 300mg

THERE'S MORE TO ASTHMA THAN YOU KNOW

WHAT YOU MAY ALREADY KNOW ABOUT SYMPTOMS:



**SHORTNESS
OF BREATH**



**CHEST
TIGHTNESS**



WHEEZING



COUGHING

BUT DID YOU KNOW ABOUT INFLAMMATION?

Inflammation leads to swelling that can cause your airways to tighten and narrow, making it harder to breathe properly. What's more, underlying inflammation can still be present, even when you don't have symptoms.

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have a parasitic (helminth) infection
- are scheduled to receive any vaccinations. You should not receive a "live vaccine" right before and during treatment with DUPIXENT.
- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
 - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

IT MAY BE TIME TO DU MORE

START WITH THE RULE OF 2 TO DETERMINE YOUR LEVEL OF ASTHMA CONTROL



Answering yes to even 1 question in the **RULE OF 2** means your asthma may not be well controlled.

- You use a rescue inhaler **2 OR MORE** times a week
- You had **2 OR MORE** courses of oral steroids in the last year
- You had **2 OR MORE** asthma attacks in the last year
- You wake up at night **2 OR MORE** times a month due to asthma

**TALK TO YOUR DOCTOR ABOUT YOUR
ASTHMA AND ASK IF DUPIXENT COULD
BE RIGHT FOR YOU.**

SEE HOW YOU MAY SAVE WITH THE
DUPIXENT MyWay[®] COPAY CARD PROGRAM ON [PAGE 16](#)

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

WHAT DUPIXENT[®] (dupilumab) CAN HELP DO

DUPIXENT requires only
1 or 2 doses a month* and can help:



**PREVENT
ASTHMA
ATTACKS**



**ACHIEVE
BETTER BREATHING
THAT LASTS**



**REDUCE OR
ELIMINATE ORAL
STEROID USE**

*Dosing based on weight and age.

UP TO
81% FEWER
ASTHMA ATTACKS^{†‡}

[†]In those with high eosinophils (a type of white blood cell).


[‡]The annual rate of severe asthma attacks was 0.20 with DUPIXENT 300 mg vs 1.04 with placebo.

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or use an asthma medicine. **Do not** change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

AND HOW IT WORKS

DUPIXENT helps block a key source of airway inflammation in the lungs.



The chronic underlying inflammation in asthma may cause airways to become constricted and narrow, making it harder to breathe.

DUPIXENT helps reduce underlying inflammation.

Illustrative image of DUPIXENT mechanism only and not reflective of treatment effect.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DU MORE YOU

In multiple clinical studies, people taking DUPIXENT saw improvements, including:

- LESS SHORTNESS OF BREATH
- LESS WHEEZING
- NOT WAKING AT NIGHT
- INCREASED ABILITY TO DO NORMAL ACTIVITIES

The relationship between DUPIXENT and these results was studied but not definitively proven.


The most common side effects include injection site reactions, pain in the throat (oropharyngeal pain), and a high count of a certain white blood cell (eosinophilia), with the addition of parasitic (helminth) infections observed in children.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including:

- **Allergic reactions. DUPIXENT can cause allergic reactions, including skin reactions, that can sometimes be severe.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye or painful red or blue bumps under the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

Some adults felt their

ASTHMA SYMPTOMS IMPROVE IN AS LITTLE AS 3 DAYS



RESULTS FOR UP TO 3 YEARS

An additional continuation study up to 3 years analyzed the safety and efficacy of 2,282 eligible patients who completed a parent study. Patients received 300 mg of DUPIXENT every 2 weeks.

In this study, all researchers knew participants received DUPIXENT, which could have influenced the results. Since participants completed a previous study, results might have favored those who already responded well to DUPIXENT. The parent trials were designed to avoid these biases.



Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

RELYING ON ORAL STEROIDS?

While they may be helpful when prescribed, long-term use of oral steroids can cause:

Potential risks of oral steroids include:

- WEIGHT GAIN
- VISION PROBLEMS
- DIABETES
- HIGH BLOOD PRESSURE
- MOOD CHANGE

This may not be an exhaustive list of side effects associated with oral steroid use.

I had to make sure my bag was packed with my daily steroids that I needed to get through the day. Since I started DUPIXENT, I've been able to decrease my reliance on oral steroids.



Tim

Real DUPIXENT patient
Individual results may vary.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including: (cont'd)

- **Inflammation of your blood vessels.** Rarely, this can happen in people with asthma who receive DUPIXENT. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. Tell your healthcare provider right away if you get: rash, chest pain, worsening shortness of breath, brown or dark colored urine, persistent fever, or a feeling of pins and needles or numbness of your arms or legs.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DUPIXENT[®] (dupilumab) IS THE ONLY BIOLOGIC SPECIFICALLY APPROVED FOR ASTHMA PATIENTS DEPENDENT ON ORAL STEROIDS



IN A STUDY OF PEOPLE DEPENDENT ON STEROIDS:

86% OF PEOPLE ON DUPIXENT
REDUCED OR ELIMINATED
THEIR ORAL STEROID USE*†

*In a clinical study of patients 12+ years old with DUPIXENT 300 mg + standard of care (SOC) vs 68% with placebo + SOC.

†Do not stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.



KIDS CAN DU MORE, TOO

Let them play! With just 1 or 2 doses a month,* DUPIXENT works continuously to help children 6+ years of age:



Breathe better in as little as 2 weeks and achieve better breathing that lasts



Reduce oral steroid use[†]



Have less asthma attacks[‡]

*Dosing based on weight and age.

[†]Do not stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.

[‡]The annual rate of severe asthma attacks was 0.24 with DUPIXENT 200 mg vs 0.67 with placebo.

The safety results of DUPIXENT observed in children through Week 52 were similar to the safety profile from studies in adults and adolescents with moderate-to-severe asthma, with the addition of parasitic (helminth) infections.

As his mom, it was so, so scary, watching Zeke go through an asthma attack. Looking back on this time, I really wish I would have taken action sooner, knowing what I know today.



Zeke, age 8

Ashley

Real DUPIXENT patient caregiver/Zeke's mom

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including: (cont'd)

- **Psoriasis.** This can happen in people with asthma who receive DUPIXENT. Tell your healthcare provider about any new skin symptoms. Your healthcare provider may send you to a dermatologist for an examination if needed.
- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

TALK TO YOUR DOCTOR

Use these questions to organize your thoughts and bring the answers to your next appointment.

START WITH THE RULE OF 2 TO DETERMINE YOUR LEVEL OF ASTHMA CONTROL



1. Do you use your rescue inhaler **2 or more** times during a typical week?

Yes No

2. Did you use **2 or more** courses of oral steroids in the last year?

Yes No

3. Did you have **2 or more** asthma attacks in the last year?

Yes No

4. Have you woken up **2 or more** nights in the past month because of your asthma?

Yes No

5. How long have you been experiencing asthma symptoms?

_____ Year(s) _____ Month(s)

6. What treatments have you or your loved one been prescribed for asthma?

Select all that apply:

Inhalers Nebulizers Biologics
 Inhaled steroids Inhaled long-acting beta antagonists
 Oral steroids None Other _____

7. How often do these symptoms occur?

Coughing

Never Rarely Occasionally Frequently

Wheezing

Never Rarely Occasionally Frequently

Rapid breathing

Never Rarely Occasionally Frequently

Chest tightness

Never Rarely Occasionally Frequently

Excess Mucus

Never Rarely Occasionally Frequently

8. Have you limited or missed an activity?

Yes No

9. Have you noticed any of the following triggers for your asthma?

Dust mites Air pollution Pets Mold
 Perfumes Scented soaps Physical exercise
 Other _____

IMPORTANT SAFETY INFORMATION: The most common side effects in patients with asthma include injection site reactions, high count of a certain white blood cell (eosinophilia), pain in the throat (oropharyngeal pain), and parasitic (helminth) infections.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

CONNECT WITH AN ASTHMA MENTOR

The **DUPIXENT Mentor Program** is designed to help new patients know what to expect by talking one-on-one with a real DUPIXENT patient mentor.

You can:

- LEARN ABOUT THEIR DUPIXENT EXPERIENCE
- ASK QUESTIONS
- GET HELPFUL TIPS



SCAN OR [CLICK HERE TO REQUEST A MENTOR.](#)

As long as you've discussed DUPIXENT with your doctor or have been prescribed DUPIXENT, you can register for a mentor so you can hear about their journey and experience with DUPIXENT.

A nurse at my doctor's office guided me through my first dose and demonstrated the injection process. After that, I felt comfortable administering my injections at home.



Bill
Real DUPIXENT patient

HOW TO TAKE DUPIXENT[®] (dupilumab)

DUPIXENT is taken by injection under the skin (subcutaneous injection). DUPIXENT should be kept in the fridge and can be removed to administer at room temperature (up to 77°F/25°C) and can be kept for up to 14 days. Remember to always review the full Instructions For Use before injecting.


It requires 1 or 2 doses per month.*

*Based on weight and age.

WHERE TO ADMINISTER DUPIXENT

Administer the subcutaneous injection into the:

- **Outer upper arms**
only if a caregiver is administering injection
- **Abdomen & Thighs**
except for the 2 inches around the navel



Choose a different site each time you inject DUPIXENT.

Do not inject into skin that is tender, damaged, bruised, or scarred.




SCAN OR [CLICK HERE FOR ADDITIONAL INJECTION SUPPORT](#)

IMPORTANT SAFETY INFORMATION: Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

SUPPORT YOU DESERVE

DUPIXENT MyWay[®] may be able to help you with the costs of your DUPIXENT treatment.



ACTIVATING YOUR COPAY CARD

You can access your Copay Card in 2 ways:



Call **1-855-726-4702**.

To ensure Copay benefits are applied to your script, make sure you have the group number and ID numbers found on the Copay Card



Visit [DUPIXENT.com](https://www.dupixent.com) to activate the card

If you are activating a Copay Card, make sure to give your Copay Card numbers to your specialty pharmacy. You can call them or share the information when they call you to schedule delivery of DUPIXENT.

The pharmacy will apply the card to lower your out-of-pocket costs with each fill of DUPIXENT, subject to the annual program maximum and other terms and conditions.

YOU COULD
PAY AS LITTLE AS **\$0***
IN COPAY FOR
DUPIXENT[®] (dupilumab)

99% OF COMMERCIALLY INSURED PATIENTS
ARE NATIONALLY COVERED FOR DUPIXENT!

If you have commercial health insurance, you may be eligible to pay **as little as \$0* in copay** per fill of DUPIXENT through the DUPIXENT MyWay Copay Card Program.

*Subject to the program maximum per patient per calendar year. Approval is not guaranteed. **THIS IS NOT INSURANCE.** Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs, including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed, or restricted. **DUPIXENT MyWay** reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.

YOU MAY BE ELIGIBLE FOR THE COPAY CARD PROGRAM IF YOU:

- Have commercial insurance, including health insurance exchanges, federal employee plans, or state employee plans
- Are prescribed DUPIXENT for an indication approved by the FDA
- Are a resident of any of the 50 United States, the District of Columbia, Puerto Rico, Guam, or the U.S. Virgin Islands

NOT ELIGIBLE?

DUPIXENT MyWay can discuss financial assistance options that may be available to you.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

STARTING *DUPIXENT MyWay*[®]

YOU CAN ENROLL IN 2 WAYS:

- Your healthcare provider can work with you to complete an enrollment form. The enrollment form must be completed and signed by you and your doctor where noted. Your doctor can then fax or upload the document electronically
- If your doctor has not already enrolled you, simply contact us directly at **1-844-DUPIXENT (1-844-387-4936) Option 1**, Monday–Friday, 8 am–9 pm ET

Your healthcare provider is your best resource for information about DUPIXENT and should be your primary point of contact throughout your treatment journey.

WHAT CAN *DUPIXENT MyWay* DO FOR YOU?

Enrolling in *DUPIXENT MyWay* can help ensure you receive DUPIXENT as quickly as possible and receive additional support along your treatment journey. For eligible patients, *DUPIXENT MyWay* can:

-  Research and review your insurance benefits with you to help you understand what your out-of-pocket costs for DUPIXENT may be.
-  Connect you with a nurse educator who can offer supplemental injection training in person, by phone, or virtually.
-  Connect you to trained DUPIXENT patient mentors to hear about their personal journeys and discuss their experiences.
-  Access to the MyWay App to help manage your DUPIXENT experience right from your mobile phone.
-  Provide sharps container for safe needle disposal.
-  Remind you to refill your prescription and take your next dose.



Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DID YOU KNOW?

DUPIXENT is also FDA-approved
in 8 other conditions:



Atopic Dermatitis
Uncontrolled
moderate-to-severe
eczema
For ages 6+ months



**Chronic Rhinosinusitis
with Nasal Polyps**
Uncontrolled
For ages 12+ years



**Eosinophilic
Esophagitis**
For ages 1+ years,
who weigh at least
33 lb (15 kg)



**Chronic Obstructive
Pulmonary Disease**
Inadequately controlled
with high blood eosinophils
For ages 18+ years



Prurigo Nodularis
For ages 18+ years

DUPIXENT is not used to
relieve sudden breathing
problems and will not replace
an inhaled rescue medicine.



**Chronic Spontaneous
Urticaria (CSU)**
with hives not controlled
by H1 antihistamines
For ages 2+ years



**Allergic fungal
rhinosinusitis**
For ages 6+ years who have
had surgery on their nose or
sinuses in the past.

DUPIXENT is not used
to treat any other forms
of hives (urticaria).



**Bullous
Pemphigoid (BP)**
For ages 18+ years



Scan or [click here](#) for more
information on what DUPIXENT
can do for you.

IMPORTANT SAFETY INFORMATION: Do not use if you are
allergic to dupilumab or to any of the ingredients in DUPIXENT[®].

Please see additional Important Safety Information throughout
and accompanying full [Prescribing Information](#) including
[Patient Information](#).

Reference:

1. IQVIA NBRx data through March 2026.

sanofi | **REGENERON**[®]

Health information contained herein is provided for general educational purposes only.
Your doctor is the single best source of information regarding your health. Please consult
your doctor if you have any questions about your health or treatment.

© 2026 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. DUPIXENT and DUPIXENT MyWay are registered trademarks of
Sanofi or an affiliate. US.DUP.25.09.0232 04/2026