



DUPIXENT[®]
(dupilumab) Injection 300mg

THE FIRST FDA-APPROVED TREATMENT FOR EoE (EOSINOPHILIC ESOPHAGITIS)

Ask your EoE specialist about DUPIXENT

INDICATION:

DUPIXENT is a prescription medicine used to treat adults and children 12 years of age and older, who weigh at least 88 pounds (40 kg), with eosinophilic esophagitis (EoE). It is not known if DUPIXENT is safe and effective in children with eosinophilic esophagitis under 12 years of age and who weigh at least 88 pounds (40 kg).

IMPORTANT SAFETY INFORMATION:

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DUPIXENT IS A BREAKTHROUGH TREATMENT FOR EoE

DUPIXENT WAS PROVEN TO REDUCE DIFFICULTY SWALLOWING*

DUPIXENT is a revolutionary biologic treatment available for patients aged 12 years and older who weigh at least 88 lb (40 kg) for EoE.

This brochure has information about DUPIXENT and can help guide you to additional resources. However, your EoE specialist is the best source of information. Be sure to ask them any questions you have before and while taking DUPIXENT.

*Based on clinical trials with 240 adult and pediatric patients (12+ years, weighing at least 88 lb) with EoE at week 24.

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you: have a parasitic (helminth) infection; are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT; are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby. A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>; are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.



DUPIXENT reduced eosinophils in the esophagus, which contribute to inflammation.*



DUPIXENT targets an underlying source of inflammation, which is a cause of EoE.



DUPIXENT is a weekly injectable treatment given under the skin (subcutaneous injection). It is not a steroid or immunosuppressant.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

QUESTIONS TO ASK YOUR DOCTOR



When talking to your doctor about managing your EoE or treatment options, you may want to discuss the following:

- Symptoms you are experiencing, such as difficulty swallowing, and their frequency
- Your current approach to managing symptoms, such as with diet and medications
- How DUPIXENT can help manage EoE symptoms



For a doctor discussion guide, scan with your smart phone.

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have EoE and asthma and use an asthma medicine. **Do not** change or stop your corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine or other asthma medicine to come back.

HOW SHOULD I USE DUPIXENT?

See the detailed “Instructions for Use” that comes with DUPIXENT for information on how to prepare and inject DUPIXENT and how to properly store and throw away (dispose of) used DUPIXENT pre-filled syringes and pre-filled pens.

Use DUPIXENT exactly as prescribed by your healthcare provider.

Your healthcare provider will tell you how much DUPIXENT to inject and how often to inject it.

DUPIXENT is taken every week as a single-dose injection using either a pre-filled syringe or a pre-filled pen.

DUPIXENT is given as an injection under the skin (subcutaneous injection).

Do not try to inject DUPIXENT until you have been shown the right way by your healthcare provider.

In children 12 years of age and older, it is recommended that DUPIXENT be given by or under the supervision of an adult.

If you miss a dose of DUPIXENT, give the DUPIXENT injection as soon as possible and start a new every week dose schedule from the time you remember to take your DUPIXENT injection.

If you inject more DUPIXENT than prescribed, get medical help or contact a Poison Center expert right away at 1-800-222-1222.

Your healthcare provider may prescribe other medicines to use with DUPIXENT. Use with other prescribed medicines exactly as your healthcare provider tells you to.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

BEGIN YOUR JOURNEY



The road to DUPIXENT begins with *DUPIXENT MyWay*. Your EoE specialist is your best resource for information about DUPIXENT and should be your primary point of contact throughout your treatment journey.

DUPIXENT MyWay can provide additional services for you and your doctor to help ensure you receive DUPIXENT as quickly as possible and receive additional support when you would like it.

SIGN UP FOR *DUPIXENT MyWay*. IT'S EASY.

Once your doctor prescribes DUPIXENT, you can enroll in the *DUPIXENT MyWay* patient support program.

Find out more at DUPIXENT.com, call **1-844-DUPIXENT (1-844-387-4936)**, **option 1**, or scan the QR code with your smart phone.



WHEN YOUR DOCTOR PRESCRIBES DUPIXENT, AND YOU ENROLL INTO *DUPIXENT MyWay*:

A Nurse Educator welcomes you

While *DUPIXENT MyWay* is confirming your insurance benefits for DUPIXENT, you'll receive a welcome call from *DUPIXENT MyWay*.

Your dedicated Nurse Educator shares helpful resources and tools from *DUPIXENT MyWay* to support you through your treatment journey.

Your prescription ships to you

Once DUPIXENT is approved, a specialty pharmacy works with you to schedule shipments of DUPIXENT to your home, or preferred location.

Your *DUPIXENT MyWay* Nurse Educator can remind you when you are due for a refill and transfer you to your selected specialty pharmacy.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

COVERING THE COST OF DUPIXENT



Before you start on DUPIXENT, DUPIXENT MyWay helps you navigate the process of accessing DUPIXENT.*

If you have insurance, *DUPIXENT MyWay*:

- Researches and helps explain your insurance benefits
- Communicates with your doctor's office about insurance approval for DUPIXENT
- Explores other potential funding sources to help with cost if your copay is unaffordable or if you are denied coverage
- The *DUPIXENT MyWay* Copay Card Program may help cover the cost of DUPIXENT for qualified commercially insured patients[†]
- If you do not have insurance, assistance may still be available
- A *DUPIXENT MyWay* Nurse Educator can help eligible patients explore additional options to help cover the cost of DUPIXENT

* Approval is not guaranteed.

[†] This offer is not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, Veterans Affairs, Department of Defense, TRICARE, or similar federal or state programs. Full program terms available at DUPIXENT.com.

NEED HELP WITH YOUR INJECTIONS?



DUPIXENT is an injection given under the skin (subcutaneous) into different injection sites. If your healthcare provider decides that you or your caregiver can do this on your own, you or your caregiver should receive training on how to prepare and inject DUPIXENT properly. **Do not** try to inject DUPIXENT until you have been shown the right way by your healthcare provider.

DUPIXENT MyWay provides supplemental injection support:

- In addition to the training from your healthcare provider, a Nurse Educator can provide supplemental injection training, either over the phone or in person
- A video about how to administer DUPIXENT is available on our website at: **DUPIXENT.com/support-savings/injection-support-center**

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF DUPIXENT?

IMPORTANT SAFETY INFORMATION

DUPIXENT can cause serious side effects, including:

- **Allergic reactions.** DUPIXENT can cause allergic reactions that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.
- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

The most common side effects in patients with eosinophilic esophagitis include injection site reactions, upper respiratory tract infections, cold sores in your mouth or on your lips, and joint pain (arthralgia).

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

ASK YOUR EoE SPECIALIST ABOUT DUPIXENT TODAY

For more detailed *DUPIXENT MyWay*[®] information, call **1-844-DUPIXENT (1-844-387-4936), option 1**, visit DUPIXENT.com/eoe or scan the QR code with your smart phone.



IMPORTANT SAFETY INFORMATION:

Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

sanofi | **REGENERON**[®]

© 2022 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. DUPIXENT[®] and *DUPIXENT MyWay*[®] are registered trademarks of Sanofi Biotechnology.
DUP.22.09.0179 10/2022

Health information contained herein is provided for general educational purposes only. Your doctor is the single best source of information regarding your health. Please consult your doctor if you have any questions about your health or treatment.

DUPIXENT[®] 
(dupilumab) Injection 300mg