

**DUPIXENT**<sup>®</sup>  
(dupilumab) Injection  
200mg • 300mg

**HELP  
HEAL  
YOUR  
SKIN  
FROM  
WITHIN**<sup>™</sup>

## STAY AHEAD OF ECZEMA WITH DUPIXENT

Available for patients as young as 6 months of age with uncontrolled moderate-to-severe eczema (atopic dermatitis)



Actual patients.  
Individual results  
may vary.

### INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 6 months of age and older with moderate-to-severe eczema (atopic dermatitis or AD) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids. It is not known if DUPIXENT is safe and effective in children with atopic dermatitis under 6 months of age.

Please see Important Safety Information throughout and click [here](#) for full [Prescribing Information](#) and [Patient Information](#).



**CRISTAL,  
ACTUAL PATIENT**  
Passion: Choreography  
*"When I stay on top of my eczema,  
I don't worry about my skin as much."*

## KNOW WHAT TO LOOK FOR

Uncontrolled moderate-to-severe eczema (atopic dermatitis) can be difficult to manage. You might do everything right and still experience symptoms. You may want to ask your doctor about making a change to your treatment.

### TALK TO AN ECZEMA SPECIALIST IF YOU OR YOUR CHILD (6+ MONTHS OF AGE):

- ▶ HIDE SKIN FROM OTHERS
- ▶ HAVE ECZEMA THAT KEEPS COMING BACK
- ▶ TAKE ORAL STEROIDS MORE THAN ONCE A YEAR
- ▶ TAKE IMMUNOSUPPRESSANTS MORE THAN ONCE A YEAR



Remember that communication is the first step towards feeling better. Scan this QR code or [CLICK HERE](#) to download our Doctor Discussion Guide and help jump-start the conversation with your doctor.

### DID YOU KNOW?

- ▶ YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

**IMPORTANT SAFETY INFORMATION:** Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®. Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you: have eye problems; have a parasitic (helminth) infection; are scheduled to receive any vaccinations. You should not receive a "live vaccine" right before and during treatment with DUPIXENT.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

# A DIFFERENCE INSIDE AND OUT

When topical prescriptions aren't enough, and your symptoms and flare-ups keep coming back, it's time for a different kind of treatment. DUPIXENT is a breakthrough treatment for uncontrolled moderate-to-severe eczema.

## DUPIXENT:

- ▶ IS A BIOLOGIC THAT HELPS BLOCK A KEY SOURCE OF INFLAMMATION INSIDE THE BODY THAT CAN CAUSE ECZEMA
- ▶ HELPS YOU STAY AHEAD OF SYMPTOMS FOR CLEARER SKIN AND LESS ITCH
- ▶ CONTINUOUSLY TREATS ECZEMA, EVEN BETWEEN FLARE-UPS
- ▶ IS NOT A STEROID OR IMMUNOSUPPRESSANT



**GRACE,  
ACTUAL PATIENT**  
Passion: Soccer

**IMPORTANT SAFETY INFORMATION:** Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you (cont'd): are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby. A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby. To enroll or get more information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>; are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see **Important Safety Information** throughout and click here for full **Prescribing Information** and **Patient Information**.



**KEVIN,  
ACTUAL PATIENT**


Passion: Volunteer Firefighter

*"Before DUPIXENT, my skin was irritated and itchy, and even worse with all my gear on."*

**OVER  
350,000\*  
PATIENTS  
PRESCRIBED DUPIXENT**

\*New adult and pediatric patients (6 months of age and older) with uncontrolled moderate-to-severe eczema.

IQVIA National Source of Business (NSOB) data through 3/17/2023



**KENNY,  
ACTUAL PATIENT**

Passion: Kung-Fu Artist

*“Before DUPIXENT, my skin was so itchy, especially when training.”*



SCAN CODE OR  
[CLICK HERE](#) TO LEARN MORE  
ABOUT HOW DUPIXENT CAN HELP.

# TAKE CONTROL TAKE YOUR SHOT AGAINST ECZEMA

Many people with moderate-to-severe eczema have an overly active immune system which produces more inflammation under the skin than usual.

DUPIXENT is a breakthrough eczema treatment. DUPIXENT works by targeting an underlying source of inflammation that may be causing your eczema symptoms. DUPIXENT works continuously on the inside to help heal your eczema on the outside.

Biologics are specialty medicines that are processed in the body differently than oral or topical medications. To be effective and work in the body properly, most biologics are delivered by injection. Biologics are used to treat a variety of conditions, including chronic inflammatory diseases that affect your immune system.



**JOLIE,  
ACTUAL PATIENT**  
Passion: Dance

**IMPORTANT SAFETY INFORMATION:** Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see [Important Safety Information](#) throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

# MEANINGFUL RESULTS FROM CLINICAL TRIALS



## HOW WERE THE TRIALS SET UP?

DUPIXENT was studied in 6 clinical trials with 2,899 patients, ages 6 months through adult, to assess its safety profile, and to determine its efficacy in clearing skin and reducing itch.

Every age group studied had moderate and/or severe eczema that was uncontrolled with topical prescription medications.

DUPIXENT was first approved to treat adults with uncontrolled, moderate-to-severe eczema by the FDA in 2017.



## WHAT DID THE TRIALS MEASURE?

The trials measured skin clearance, skin improvement, itch reduction and safety among people taking DUPIXENT compared to those who were not taking DUPIXENT. Each trial measured the results at 16 weeks. One trial measured the results at 16 and 52 weeks (1 year).



## WHAT MEDICATIONS WERE USED?

Depending on the trial, participants received one of the following treatments:

- ▶ DUPIXENT
- ▶ Placebo\*
- ▶ DUPIXENT with a topical corticosteroid (TCS)
- ▶ Placebo with TCS

\*A treatment that appears to be real but contains no medicine.

## DID YOU KNOW?

- ▶ YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

**IMPORTANT SAFETY INFORMATION:** Especially tell your healthcare provider if you are taking oral, topical or inhaled corticosteroid medicines or if you have atopic dermatitis and asthma and use an asthma medicine.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**LEI ANN,  
ACTUAL PATIENT**

Passion: Traveling

*"It wasn't always easy traveling with my eczema. Now, with DUPIXENT, I don't worry as much about my skin."*

# HEALING FROM WITHIN IS A POWERFUL THING

**MOSETTA,  
ACTUAL PATIENT**

Passion: Fitness Enthusiast

“Before DUPIXENT,  
eczema made my skin so  
uncomfortable it was  
tough working out.”

TRIAL RESULTS: DUPIXENT vs PLACEBO\*

Two separate 16-week clinical trials:

**~4X SAW CLEAR OR ALMOST CLEAR SKIN**

37% vs 9% not taking DUPIXENT (placebo)

**>3X HAD SIGNIFICANT ITCH REDUCTION**

38% vs 11% not taking DUPIXENT

ALMOST HALF OF ADULTS SAW

**≥75% SKIN IMPROVEMENT**

48% vs 13% not taking DUPIXENT

A PORTION EVEN SAW

**≥90% SKIN IMPROVEMENT**

33% vs 7% not taking DUPIXENT

**SAFETY RESULTS:**

The most common side effects observed in clinical trials were injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, cold sores in the mouth or on the lips, and high count of a certain white blood cell (eosinophilia).

Long-term safety results for adults continuing on DUPIXENT in the extension study for nearly 3 years were consistent with the adult clinical trials.

\*A treatment that appears to be real but contains no medicine.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

SCAN CODE OR  
CLICK HERE TO HEAR  
ACTUAL PATIENT STORIES.



ADULTS (AGES 18+)

# LASTING CHANGE IS ACHIEVABLE

TRIAL RESULTS: DUPIXENT + TOPICAL CORTICOSTEROIDS (TCS) vs TCS ONLY

Long-term results from a clinical trial that studied  
DUPIXENT for 52 weeks.

## CLEAR OR ALMOST CLEAR SKIN AT 16 WEEKS

39% taking DUPIXENT + TCS vs 12% using TCS only

## LONG-LASTING CLEARER SKIN AT 16 AND 52 WEEKS

22% taking DUPIXENT + TCS vs 7% using TCS only

## FAST ITCH RELIEF AS SOON AS 2 WEEKS

18% taking DUPIXENT + TCS vs 8% using TCS only

## SIGNIFICANT ITCH REDUCTION AT 16 WEEKS

59% taking DUPIXENT + TCS vs 20% using TCS only

## ITCH RELIEF THAT LASTS AT 52 WEEKS

51% taking DUPIXENT vs 13% using TCS only

**IMPORTANT SAFETY INFORMATION:** Do not change or stop your corticosteroid medicine or other asthma medicine without talking to your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine or other asthma medicine to come back.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**KEVIN,  
ACTUAL PATIENT**  
Passion: Volunteer Firefighter

## GET AHEAD WITH CLEARER SKIN AND LESS ITCH

TRIAL RESULTS: DUPIXENT vs PLACEBO\*

Teens took part in a 16-week clinical trial.

### CLEARER SKIN

# 12X SAW CLEAR OR ALMOST CLEAR SKIN

24% vs 2% not taking DUPIXENT (placebo)

### NOTICEABLY LESS ITCH

# ~7X EXPERIENCED ITCH RELIEF

37% vs 5% not taking DUPIXENT

*\*A treatment that appears to be real but contains no medicine.*

**IMPORTANT SAFETY INFORMATION:** DUPIXENT can cause serious side effects, including: Allergic reactions. DUPIXENT can cause allergic reactions that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, joint pain, general ill feeling, itching, skin rash, swollen lymph nodes, nausea or vomiting, or cramps in your stomach-area.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

### JOLIE, ACTUAL PATIENT

Passion: Dance

"My skin was so itchy, especially with my dance outfit. But now I notice a difference."

TEENS (AGES 12-17)

# PROVEN RESULTS AND DEMONSTRATED SAFETY PROFILE IN TEENS

TRIAL RESULTS CONTINUED: DUPIXENT vs PLACEBO\*

Teens took part in a 16-week clinical trial.

NEARLY HALF OF TEENS SAW

**≥75% SKIN IMPROVEMENT**

42% vs 8% in those not taking DUPIXENT

SOME EVEN SAW

**≥90% SKIN IMPROVEMENT**

23% vs 2% not taking DUPIXENT

► **YOU CAN TAKE DUPIXENT WITH OR WITHOUT  
A TOPICAL CORTICOSTEROID (TCS)**

#### SAFETY RESULTS:

Long-term safety results for teens continuing on DUPIXENT in the pediatric extension study through 1 year were consistent with the adult clinical trials.

\*A treatment that appears to be real but contains no medicine.

**IMPORTANT SAFETY INFORMATION:** DUPIXENT can cause serious side effects, including (cont'd): Eye problems. Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**TYLER,  
ACTUAL PATIENT**

Passion: Kayaking

*"Before DUPIXENT, I was always so itchy. Especially when I was hot."*

## RESULTS THEY CAN SEE RESULTS THEY CAN FEEL

Caring for your child with uncontrolled moderate-to-severe eczema can be a struggle—especially when you're constantly trying to avoid triggers.

### TRIAL RESULTS: DUPIXENT + TCS vs TCS ONLY

Children took part in a 16-week clinical trial.

### CLEARER SKIN

**>2X SAW CLEAR OR ALMOST CLEAR SKIN**

30% on 300 mg every 4 weeks + TCS vs 13% on TCS only  
39% on 200 mg every 2 weeks + TCS vs 10% on TCS only

### NOTICEABLY LESS ITCH

**~4.5X HAD LESS ITCH**

54% on 300 mg every 4 weeks + TCS vs 12% on TCS only  
61% on 200 mg every 2 weeks + TCS vs 13% on TCS only

**IMPORTANT SAFETY INFORMATION:** DUPIXENT can cause serious side effects, including (cont'd): Eye problems. Your healthcare provider may send you to an ophthalmologist for an eye exam if needed.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

**HARPER,  
ACTUAL PATIENT**  
Passion: Running

# PROVEN RESULTS AND DEMONSTRATED SAFETY PROFILE IN CHILDREN

TRIAL RESULTS CONTINUED: DUPIXENT + TCS vs TCS ONLY

Children took part in a 16-week clinical trial.

MORE THAN HALF OF CHILDREN SAW

**≥75% SKIN IMPROVEMENT**

75% on 300 mg every 4 weeks + TCS vs 28% on TCS only

75% on 200 mg every 2 weeks + TCS vs 26% on TCS only

**YOUR HEALTHCARE PROVIDER WILL TELL YOU HOW MUCH  
DUPIXENT TO INJECT AND HOW OFTEN.**

► **YOU CAN TAKE DUPIXENT WITH OR WITHOUT  
A TOPICAL CORTICOSTEROID (TCS)**

**SAFETY RESULTS:**

Long-term safety results for children continuing on DUPIXENT with or without TCS in the pediatric extension study through 1 year were consistent with the adult clinical trials.

**IMPORTANT SAFETY INFORMATION:** DUPIXENT can cause serious side effects, including (cont'd): Joint aches and pain. Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).



**GRACE,  
ACTUAL PATIENT**  
Passion: Soccer

## HEALING FROM WITHIN FOR YOUR LITTLE ONE

DUPIXENT is the only FDA-approved biologic for treating uncontrolled moderate-to-severe eczema in children as young as 6 months.

TRIAL RESULTS: DUPIXENT + TCS vs TCS ONLY

Young children took part in a 16-week clinical trial.

### CLEARER SKIN 7X SAW CLEAR OR ALMOST CLEAR SKIN

28% on DUPIXENT + TCS vs 4% on TCS only

### ITCH RELIEF ~ 5X HAD LESS ITCH

48% on DUPIXENT + TCS vs 9% on TCS only

### PROVEN RESULTS HALF OF YOUNG CHILDREN SAW ≥75% SKIN IMPROVEMENT

53% on DUPIXENT + TCS vs 11% on TCS only

#### ▶ YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

#### SAFETY RESULTS:

Long-term safety results in young children continuing on DUPIXENT with or without TCS in the pediatric extension study through 1 year were consistent with the adult clinical trials. In addition, hand-foot-and-mouth disease and warts were reported. In these reported cases, patients continued treatment with DUPIXENT.

**IMPORTANT SAFETY INFORMATION:** DUPIXENT can cause serious side effects, including: Joint aches and pain (cont'd). Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

Not an actual patient.

# HELPS HEAL THE LOOK AND FEEL OF SKIN

Eczema can affect anyone, regardless of race, ethnicity, or gender, and it can appear differently on different skin types. In some cases, eczema rashes may look red, gray, brown or purple. DUPIXENT has shown generally consistent efficacy regardless of race/ethnicity in adult studies.



Actual 4-year-old patient in the clinical trial.† Individual results may vary.



Actual adult patient.\*



Actual 6-year-old patient in the clinical trial.† Individual results may vary.

**IMPORTANT SAFETY INFORMATION:** The most common side effects in patients with eczema include injection site reactions, eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, cold sores in your mouth or on your lips, and high count of a certain white blood cell (eosinophilia).

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Please see **Important Safety Information** throughout and click [here for full Prescribing Information and Patient Information](#).

\*This adult patient was an actual patient treated with DUPIXENT. Not clinical trial patient. Scoring was designated by the treating physician. Because this was a real-world patient, other factors may have influenced their treatment results. Individual results may vary.

† All patients were prescribed low or medium-potency topical corticosteroids (TCS) based on the clinical trial program.

## DOSING FOR DUPIXENT® (dupilumab):

	BODY WEIGHT	INITIAL LOADING DOSE	SUBSEQUENT DOSAGE
<b>ADULTS 18+</b>	Any weight	600 mg (two 300-mg injections)	300 mg; every 2 weeks
<b>CHILDREN &amp; TEENS 6 TO 17 YEARS OF AGE</b>	132 lb or more	600 mg (two 300-mg injections)	300 mg; every 2 weeks
	66 lb to under 132 lb	400 mg (two 200-mg injections)	200 mg; every 2 weeks
	33 lb to under 66 lb	600 mg (two 300-mg injections)	300 mg; every 4 weeks
	BODY WEIGHT	INITIAL* AND SUBSEQUENT DOSAGE	
<b>CHILDREN 6 MONTHS TO 5 YEARS OF AGE</b>	33 lb to under 66 lb	300 mg; every 4 weeks	
	11 lb to under 33 lb	200 mg; every 4 weeks	

\*For children 6 months to 5 years of age with atopic dermatitis, no initial loading dose is recommended.

### DUPIXENT is available as a pre-filled pen or pre-filled syringe.

Your doctor will tell you how much DUPIXENT to inject and how often to inject it, following proper training. DUPIXENT is an injection given under the skin (subcutaneous injection). Please rotate the injection site every time you administer DUPIXENT.

DUPIXENT can be given by a doctor in the office or a caregiver at home:

- ▶ in children 6 months to less than 12 years, DUPIXENT should be given by a caregiver
- ▶ in teens 12-17 years, DUPIXENT should be given by or supervised by an adult



Pre-filled pen: available in 200 mg and 300 mg doses for patients aged 2+ years



SCAN CODE OR [CLICK HERE TO WATCH THE TIPS AND TRICKS VIDEO ON HOW TO ADMINISTER DUPIXENT](#)



Pre-filled syringe: available in 200 mg and 300 mg doses for patients aged 6+ months

**IMPORTANT SAFETY INFORMATION:** Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT.

Please see [Important Safety Information](#) throughout and [click here for full Prescribing Information and Patient Information](#).

# DUPIXENT myway

## WITH YOU EVERY STEP OF THE WAY

Once you've been prescribed DUPIXENT, the *DUPIXENT MyWay*® patient support program is here to help answer questions you may have about DUPIXENT. Keep in mind your doctor is still the best resource for you if you have any questions about your eczema or your treatment program.

To enroll in *DUPIXENT MyWay*® and access helpful resources, call 1-844-DUPIXENT (1-844-387-4936) and select Option 1.



**CHLOE,  
ACTUAL PATIENT**  
Passion: Surfing

## YOU'RE NOT ALONE

It's important to know that *DUPIXENT MyWay*® helps people just like you who are on or about to start DUPIXENT.



For eligible patients, enrolling in *DUPIXENT MyWay* can help ensure you receive DUPIXENT as quickly as possible and receive additional support along your treatment journey.



The *DUPIXENT MyWay Support Team* can answer questions about DUPIXENT, arrange supplemental injection training, and help you start and stay on track.



With the *DUPIXENT MyWay Patient App* powered by Medisafe, people just like you can get access to tools to help you start and stay on track with your treatment. Contact your **Case Manager** for an invitation with directions on how to get the app.



The *DUPIXENT MyWay Mentor Program* gives you the opportunity to speak with other patients and caregivers to hear their experiences.



SCAN OR CLICK HERE TO LEARN ABOUT ADDITIONAL *DUPIXENT MYWAY* RESOURCES.

**IMPORTANT SAFETY INFORMATION:** Do not try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 months to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see **Important Safety Information** throughout and click here for full **Prescribing Information** and **Patient Information**.

# DUPIXENT myway

## NEED HELP PAYING?

The DUPIXENT MyWay Copay Card Program may help reduce the out-of-pocket cost of DUPIXENT. Eligible patients may pay as little as \$0\* per fill for each DUPIXENT prescription (maximum of \$13,000 per patient per calendar year).\*

### AS LITTLE AS \$0\* COPAY MAY BE AVAILABLE IF YOU:

- ▶ Have commercial insurance
- ▶ Have a DUPIXENT prescription for an FDA-approved condition
- ▶ Reside in the 50 United States, the District of Columbia, Puerto Rico, Guam, or the USVI
- ▶ Are a patient or caregiver 18 years or older

*\*Approval is not guaranteed. Program has an annual maximum of \$13,000. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed or restricted. DUPIXENT MyWay reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.*

## OTHER OPTIONS FOR SUPPORT

Through the Patient Assistance Program, patients who are uninsured or rendered uninsured by their insurance provider could receive DUPIXENT free of charge if they qualify.

Those who may qualify must be at least 18 years of age or older, a resident of the 50 United States, the District of Columbia, or Puerto Rico, and demonstrate a financial need.

Call DUPIXENT MyWay at 1-844-387-4936 for full program information and eligibility criteria.



**DUPIXENT**   
(dupilumab) Injection  
200mg · 300mg

# HELP HEAL YOUR SKIN FROM WITHIN™

ASK YOUR ECZEMA SPECIALIST  
ABOUT DUPIXENT AND  
VISIT [DUPIXENT.COM](https://www.dupixent.com)  
OR CALL 1-844-DUPIXENT  
(1-844-387-4936)



SCAN CODE OR  
[CLICK HERE TO](#)  
LEARN MORE  
ABOUT DUPIXENT

**TYLER,**  
**ACTUAL PATIENT**  
Passion: Kayaking

**IMPORTANT SAFETY INFORMATION:** Do not use if you are allergic to dupilumab or to any of the ingredients in **DUPIXENT**®.

Please see Important Safety Information throughout and click here for full [Prescribing Information](#) and [Patient Information](#).

**sanofi** | **REGENERON**®

© 2023 Sanofi and Regeneron Pharmaceuticals, Inc. All rights reserved.  
DUPIXENT® and DUPIXENT MyWay® are registered trademarks of  
Sanofi Biotechnology. DUP.23.07.0008 07/23