

**DUPIXENT**<sup>®</sup>  
(dupilumab) Injection  
200mg · 300mg

**FIRST AND ONLY FDA-APPROVED  
BIOLOGIC FOR APPROPRIATE  
CSU PATIENTS AS YOUNG AS  
2 YEARS OLD**

**DISCOVER A  
DIFFERENT WAY  
TO TREAT CHRONIC  
SPONTANEOUS  
URTICARIA (CSU)  
ALSO KNOWN AS CHRONIC  
HIVES WITH NO KNOWN CAUSE**

DUPIXENT is a prescription medicine used to treat adults and children 2 years of age and older with chronic spontaneous urticaria (CSU), who continue to have hives that are not controlled with H1 antihistamine treatment. DUPIXENT is not used to treat any other forms of hives (urticaria).

**HELP  
HEAL  
YOUR  
SKIN  
FROM  
WITHIN<sup>™</sup>**

**TIRED OF  
PLAYING  
HIVES  
AND  
SEEK?**

Learn about a treatment option for **Chronic Spontaneous Urticaria (CSU)**, also known as chronic hives with no known cause.

Patient portrayal

#### INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 2 years of age and older with chronic spontaneous urticaria (CSU), who continue to have hives that are not controlled with H1 antihistamine treatment. **DUPIXENT is not used** to treat any other forms of hives (urticaria).

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

#### IMPORTANT SAFETY INFORMATION:

**Do not use** if you are allergic to dupilumab or to any of the ingredients in DUPIXENT<sup>®</sup>.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

**Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:**

- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

# TAKE YOUR NEXT STEP

If your antihistamine isn't doing enough to relieve your chronic spontaneous urticaria (CSU), also known as chronic hives with no known cause:

Take a moment to reflect on why you or your loved one are considering something different.

## PLEASE LIST YOUR GOALS FOR TREATMENT BELOW

---

---

---

---

---

---

---

---

---

---

---

---

### IMPORTANT SAFETY INFORMATION:

Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you (cont'd):

- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
  - o A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby.



## ASK YOUR SPECIALIST THE FOLLOWING QUESTIONS

1. How effective is DUPIXENT at treating CSU?
2. How is DUPIXENT administered?
3. When could I see results on DUPIXENT?
4. What are the benefits and side effects of DUPIXENT?
5. How do I get started with DUPIXENT?
6. What resources and support are available?

Bring this guide to your next appointment to see if DUPIXENT is an option for you.

- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.



# STUCK IN A CYCLE OF CHRONIC HIVES THAT APPEAR OUT OF NOWHERE?

These itchy red or skin-colored bumps with no known cause can ruin your day. You've tried so many treatments, but hives still show up unannounced, without rhyme or reason.

CSU is not an allergic reaction. It may be that the body's immune system is out of balance—and that could be the source of spontaneous flare-ups. This immune imbalance can be a driver of inflammation that contributes to the itch and hives you experience. CSU can be diagnosed based on your symptoms and medical history. Allergy testing and/or skin biopsy is not required for diagnosis.

If you're tired and frustrated with the relentless itch and discomfort, you're not alone.

Too often, the uncomfortable red or skin-colored bumps and swelling becomes an unpredictable cycle that happens again and again. And that can be challenging for anyone who's coping with it.



Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.



Patient portrayal

# SYMPTOMS OF CSU

## DO THESE SOUND FAMILIAR?



**ITCHY HIVES HAPPEN SPONTANEOUSLY WITH NO KNOWN CAUSES**



**6+ WEEKS OF HIVES ON AND OFF, WITH OR WITHOUT ANGIOEDEMA (DEEP SWELLING UNDER THE SKIN)**



**FLARE-UPS THAT CAN LAST FOR DAYS OR EVEN WEEKS**



**HIVES THAT APPEAR, DISAPPEAR, THEN APPEAR ON ANOTHER AREA**



**HIVES THAT OFTEN SHOW UP ON THE CHEST, BACK, ARMS, OR LEGS**

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

# MANY PEOPLE WITH CSU HAVE BEEN SEARCHING FOR ANOTHER OPTION



## DID YOU KNOW?

FOR ABOUT  
**50%**  
OF ADULT  
PATIENTS

**ANTIHISTAMINES ARE NOT ENOUGH, EVEN WHEN TAKING MORE THAN THE STANDARD DOSE**

**2-3**  
YEARS

**IS THE AVERAGE TIME ADULTS SPEND SEEKING A DIAGNOSIS**

ALMOST  
**2X**  
AS MANY

**ADULT WOMEN ARE AFFECTED BY CSU COMPARED TO MEN**

**2-5**  
YEARS

**IS THE AVERAGE LENGTH FOR MOST HIVES CASES IN ADULTS, BUT SOME CAN LAST LONGER**

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).


# SEEKING HELP WITH CSU? START HERE.

DUPIXENT is a different way to treat CSU. It's a biologic that targets a key source of inflammation that can cause symptoms and works to help address that inflammation. DUPIXENT can help you get ahead of your unpredictable symptoms by reducing the intense itch and number of hives.

What's more, when used as prescribed, DUPIXENT continuously works within your body between doses to help manage CSU symptoms.

Based on the prescribing information, DUPIXENT does not require initial lab testing or ongoing lab monitoring.

**Get ahead of symptoms with DUPIXENT.**



Patient portrayal

## **IMPORTANT SAFETY INFORMATION:**

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

# UNDERSTAND HOW IT WORKS

## **DUPIXENT:**

- ▶ **TARGETS A KEY SOURCE OF INFLAMMATION**
- ▶ **IS NOT AN IMMUNOSUPPRESSANT OR STEROID**
- ▶ **CONTINUOUSLY WORKS BETWEEN DOSES TO HELP TREAT CSU**
- ▶ **HELPS MANAGE YOUR SYMPTOMS PROACTIVELY**

## **TALK TO YOUR SPECIALIST IF YOU HAVE:**

- ▶ **SPONTANEOUS HIVES FOR 6+ WEEKS**
- ▶ **RED OR SKIN-COLORED ITCHY WELTS OR SWELLING**
- ▶ **UNPREDICTABLE FLARE-UPS**
- ▶ **TRIED ANTIHISTAMINES**

There is no specific test for CSU. Talk to your doctor about your symptoms.

# A DIFFERENCE YOU CAN SEE AND FEEL

DUPIXENT was studied in two 24-week clinical trials in 284 patients ages 12 years and older with CSU who continued to have hives despite taking H1 antihistamines. All patients remained on antihistamines during the trials.

## AT 24 WEEKS PATIENTS HAD:



### LESS ITCH\*

Trial 1: 64% less itch on DUPIXENT vs 35% on placebo  
Trial 2: 48% less itch on DUPIXENT vs 33% on placebo



### REDUCTION IN BOTH ITCH AND HIVES†

Trial 1: 66% reduction on DUPIXENT vs 36% on placebo  
Trial 2: 48% reduction on DUPIXENT vs 32% on placebo

## RELIEF MAY BE POSSIBLE:



### ~2X AS MANY PATIENTS HAD NO ITCH AND NO HIVES AT 24 WEEKS

Trial 1: 32% on DUPIXENT vs 13% on placebo  
Trial 2: 30% on DUPIXENT vs 17% on placebo

\*Patients rated itch daily (from 0-3) for 7 days (max score 21), compared to baseline score.

†Patients also rated their hives daily (from 0-3) for 7 days (max score 21). This score was added to their itch severity score, compared to baseline score.

**The most common side effects include injection site reactions.** DUPIXENT's use for treating CSU in children ages 2 - 11 years is supported by the results from two studies that included patients 12 years of age and older with CSU.

### IMPORTANT SAFETY INFORMATION:

Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have CSU and asthma and use an asthma medicine.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

# TAKING DUPIXENT<sup>®</sup> (dupilumab)

## DUPIXENT IS AVAILABLE AS A PRE-FILLED PEN OR PRE-FILLED SYRINGE

- ▶ DUPIXENT is a biologic medicine given under the skin (subcutaneous injection).
- ▶ It is not taken every day. DUPIXENT is taken only once or twice a month (based on age and weight).
  - ▶ Your doctor will tell you how much DUPIXENT to take.
- ▶ DUPIXENT can be given by your doctor or self-administered at home.
  - ▶ Do not try to prepare and inject DUPIXENT until being properly trained by your doctor.
  - ▶ In children 12 years of age and older, it is recommended that DUPIXENT be given by or under the supervision of an adult.
  - ▶ In children 2 years to less than 12 years of age, DUPIXENT should be given by a caregiver.



Click here to learn more about how to inject DUPIXENT.

### IMPORTANT SAFETY INFORMATION:

Do not change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.


## YOUR SPECIALIST WILL PRESCRIBE THE APPROPRIATE DOSAGE BASED ON THE RECOMMENDATIONS BELOW:

	BODY WEIGHT	INITIAL LOADING DOSE	SUBSEQUENT DOSAGE
<b>ADULTS 18+ YEARS</b>	Any weight	600 mg (two 300-mg injections)*	300 mg; every 2 weeks
<b>CHILDREN AND TEENS (6-17 YEARS OF AGE)</b>	132 lb or more	600 mg (two 300-mg injections)*	300 mg; every 2 weeks
	66 lb to under 132 lb	400 mg (two 200-mg injections)*	200 mg; every 2 weeks
	33 lb to under 66 lb	600 mg (two 300-mg injections)*	300 mg; every 4 weeks

\*Administered into different injection sites.

	BODY WEIGHT	RECOMMENDED DOSAGE <sup>†</sup>
<b>YOUNG CHILDREN (2-5 YEARS OF AGE)</b>	33 lb to under 66 lb	300 mg (one 300-mg injection); every 4 weeks
	11 lb to under 33 lb	200 mg (one 200-mg injection); every 4 weeks

<sup>†</sup>For pediatric patients 2 years to 5 years of age with CSU, no initial loading dose is recommended.



# HELPFUL RESOURCES FOR YOU, EVERY STEP OF THE WAY

DUPIXENT MyWay is a patient support program designed to help you start DUPIXENT as quickly as possible and stay on track with your treatment.

Of course, your doctor is always the first and best resource for any questions about CSU or your treatment program. DUPIXENT MyWay is also here to support you.

Sometimes, it's nice to know that when you need answers to your questions, you have someone by your side. A DUPIXENT MyWay Case Manager will work closely with you and your doctor to provide you with the tools and support you need.

When you enroll, a dedicated Case Manager can:



Guide you through the insurance process to help you understand your coverage options



Copay and financial assistance for eligible patients



Connect you with a Nurse Educator, who can provide supplemental injection training, and answer questions about your condition and treatment

The MyWay App can help you manage your DUPIXENT experience right from your mobile device. It offers resources to help you start and stay on track with your treatment. Ask your DUPIXENT MyWay Case Manager for access.



CLICK HERE TO LEARN MORE ABOUT DUPIXENT.

Sign up for DUPIXENT MyWay once you've been prescribed DUPIXENT. Enroll at [DUPIXENT.com](http://DUPIXENT.com) or call 1-844-DUPIXENT (1-844-387-4936).

# NEED HELP PAYING? AS LITTLE AS \$0\* COPAY MAY BE AVAILABLE



DUPIXENT MyWay may be able to help you with the costs of your prescription.

If you have commercial health insurance, you may be eligible to pay as little as \$0\* in copay per fill of DUPIXENT through the DUPIXENT MyWay Copay Card Program.

## YOU MAY BE ELIGIBLE IF YOU:

- ▶ Have commercial insurance
- ▶ Have a DUPIXENT prescription for an FDA-approved condition
- ▶ Reside in the 50 United States, the District of Columbia, or Puerto Rico
- ▶ Are a patient or caregiver 18 years or older

## THERE ARE OTHER OPTIONS FOR SUPPORT

Through the Patient Assistance Program, patients who are uninsured or rendered uninsured by their insurance provider could receive DUPIXENT at no cost, if eligible. Those who may qualify must be at least 18 years of age or older, a resident of the 50 United States, the District of Columbia, or Puerto Rico, and demonstrate a financial need. DUPIXENT MyWay reviews your situation individually and evaluates eligibility on a case-by-case basis. Find out if you qualify by speaking with a DUPIXENT MyWay Case Manager at 1-844-DUPIXENT (1-844-387-4936) Option 1.

*\*Subject to the program maximum per patient per calendar year. Approval is not guaranteed. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs, including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed, or restricted. DUPIXENT MyWay reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.*

## IMPORTANT SAFETY INFORMATION:

**DUPIXENT can cause serious side effects, including:**

- **Allergic reactions. DUPIXENT can cause allergic reactions, including skin reactions, that can sometimes be severe.** Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye or painful red or blue bumps under the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

# READY TO TALK TO YOUR SPECIALIST?

Answer these questions and bring them to your visit to have a more productive conversation.

**How long have you or your loved one been experiencing symptoms of chronic hives?**

≥6 Months    ≥2 Years    ≥5 Years    Other \_\_\_\_\_

**How often do you notice hives?**

Daily    Every other day    Weekly    Other \_\_\_\_\_

**In the last week, about how many hives have you noticed each day?**

Few (0-5)    Some (6-19)    Many (20+)

**Where are the hives usually located?** (Check all that apply)

Shoulders    Arms    Hands    Chest    Face  
 Stomach    Back    Legs    Feet    Other \_\_\_\_\_

**Rate the itchiness of the hives?**

No itch  
 Mild: Itchy but not bothersome  
 Moderate: Itch that is bothersome but not disruptive  
 Intense: Itch that is extremely disruptive as well as bothersome

**Describe any additional impact living with chronic hives has had on you or your loved one.** (Eg, My son's flare-ups are so bothersome. He often worries about how he will manage them.)

---



---

## IMPORTANT SAFETY INFORMATION:

**DUPIXENT can cause serious side effects, including (cont'd):**

- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

## WHAT'S WORKING? WHAT'S NOT?

Let your doctor know about everything you or your loved one have used to manage chronic hives.

**Have you or your loved one tried any of the following? Check all that apply, and note frequency of use.**

	Daily	Weekly	Biweekly	Monthly	A few times a year
Antihistamines	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Steroids	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Immunosuppressants	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Other prescription medicines	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
_____					
Over-the-counter medicines	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
_____					
Other therapies (eg, supplements)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

**How well does your current treatment plan provide symptom relief? If you still have bothersome symptoms despite treatment, please list them below.**

---



---



---



**Tip:** Take photos of your flare-ups to show your doctor and discuss at your visit.

**DUPIXENT**  
(dupilumab) Injection  
200mg · 300mg

**HELP  
HEAL  
YOUR  
SKIN  
FROM  
WITHIN™**



Patient portrayal

## CONTROLLING CHRONIC HIVES WITH NO KNOWN CAUSE IS POSSIBLE

ASK YOUR SPECIALIST ABOUT DUPIXENT AND VISIT  
DUPIXENT.COM OR CALL 1-844-DUPIXENT (1-844-387-4936)

### IMPORTANT SAFETY INFORMATION:

The most common side effects in patients with chronic spontaneous urticaria include injection site reactions.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 2 years to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

**sanofi** | **REGENERON**®

© 2026 Sanofi and Regeneron Pharmaceuticals, Inc. All rights reserved. DUPIXENT® and DUPIXENT MyWay® are registered trademarks of Sanofi Biotechnology. US.DUP.26.04.0298 5/26



[CLICK HERE TO LEARN MORE  
ABOUT DUPIXENT.](#)