

DUPIXENT[®]
(dupilumab) Injection
200mg · 300mg

DUPIXENT HELPS YOU
**FEEL
THE HEAL**
AND SEE THE DIFFERENCE

FDA-APPROVED FOR AGES 6 MONTHS AND UP



Actual patient. Individual results may vary.

THE #1 PRESCRIBED BIOLOGIC FOR MODERATE-TO-SEVERE ECZEMA

INDICATION

DUPIXENT is a prescription medicine used to treat adults and children 6 months of age and older with moderate-to-severe eczema (atopic dermatitis or AD) that is not well controlled with prescription therapies used on the skin (topical), or who cannot use topical therapies. DUPIXENT can be used with or without topical corticosteroids. It is not known if DUPIXENT is safe and effective in children with AD under 6 months of age.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

WHEN IS IT TIME TO RETHINK YOUR ECZEMA TREATMENT?

Eczema can be challenging, even when you're already following your doctor's advice and using topical prescription (Rx) creams and ointments to manage it. You may still struggle with dry, itchy rashes and continuing flare-ups. It's a situation many people with moderate-to-severe eczema deal with, because sometimes topical Rx treatments aren't enough. The itchy rashes keep recurring due in part to the inflammation below the skin's surface. This inflammation is caused by an overactive immune system.

Maybe it's time to ask your doctor for a treatment that works differently.

TALK TO AN ECZEMA SPECIALIST WHEN YOU OR YOUR CHILD:

- ▶ KEEP USING TOPICAL Rx CREAMS AND OINTMENTS THAT AREN'T CONTROLLING YOUR ECZEMA
- ▶ FREQUENTLY HAVE ITCHY, UNCOMFORTABLE SKIN AND FLARE-UPS
- ▶ TAKE ORAL STEROIDS OR IMMUNOSUPPRESSANTS MORE THAN ONCE A YEAR
- ▶ HIDE YOUR SKIN FROM OTHERS



[CLICK HERE](#) TO PREP FOR A
CONVERSATION WITH YOUR DOCTOR



**KATE,
ACTUAL PATIENT**

IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

INFLAMMATION INSIDE CAN CAUSE ECZEMA OUTSIDE

You may be treating your moderate-to-severe eczema with just topical Rx creams and ointments. But is it enough if your itchy rashes keep coming back?

Even though eczema appears on your skin, the flare-ups and irritating itch are due in part from inflammation underneath your skin. **That's why it's worth exploring a treatment that works from the inside, like DUPIXENT.**

If you have eczema, even the simplest touch can be uncomfortable. Everyday interactions, from handshakes to hugs, can be tough on your rashy skin.

DUPIXENT can help you feel the heal and see the difference with less itch and clearer skin.



CAMERON,
ACTUAL PATIENT

DUPIXENT[®] (dupilumab) WORKS CONTINUOUSLY TO:

- ▶ HELP BLOCK A KEY SOURCE OF INFLAMMATION INSIDE THE BODY THAT CAN CAUSE ECZEMA
- ▶ HELP YOU STAY AHEAD OF ECZEMA WITH LESS ITCH AND CLEARER SKIN
- ▶ TREAT ECZEMA, EVEN BETWEEN FLARE-UPS

DID YOU KNOW? DUPIXENT:

- ▶ IS NOT A STEROID OR IMMUNOSUPPRESSANT
- ▶ IS NOT TAKEN DAILY
- ▶ CAN BE TAKEN WITH OR WITHOUT A TOPICAL CORTICOSTEROID

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you:

- have eye problems.
- have a parasitic (helminth) infection.
- are scheduled to receive any vaccinations. You should not receive a “live vaccine” right before and during treatment with DUPIXENT.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).


CONSIDER A DIFFERENT APPROACH

DUPIXENT is a biologic. Biologics are specialty medicines used to treat a variety of conditions, including chronic inflammatory diseases such as moderate-to-severe eczema.

Biologics are processed in the body differently than oral or topical medications. To be effective and work properly, most biologics are injectable medicines.

If you've been struggling to manage your eczema with just prescription topicals, it may be time to consider a biologic like DUPIXENT. It helps block a key source of inflammation under the skin.

DUPIXENT can help you feel the heal and see the difference with less itch and clearer skin.



*New adult and pediatric patients (6 months of age and older) with uncontrolled moderate-to-severe eczema. IQVIA National Source of Business (NSOB) data through March 2025.

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you (cont'd):

- are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby.
 - A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby.
- are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.



Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.



KENNY,
ACTUAL PATIENT

WHAT IS A CLINICAL TRIAL?

Before a drug is made available to the general public, its efficacy and safety need to be demonstrated through controlled clinical trials.* After completing these trials, the information is sent to the FDA for their review. The FDA reviews the trial results and makes a decision on whether to approve the drug.

WHO WAS INCLUDED IN THE TRIALS?

DUPIXENT was studied in 7 eczema clinical trials with 3,032 patients, ages 6 months and older.

Every age group had moderate and/or severe eczema that was not well controlled.

WHAT DID THE TRIALS MEASURE?

The trials measured skin clearance, itch reduction, and safety among people taking DUPIXENT compared to those not taking it. Each trial measured results at 4 months (16 weeks). One of the trials also measured results at 1 year (52 weeks).

WHAT MEDICATIONS WERE USED?

Depending on the trial, participants received one of the following:

- ▶ DUPIXENT
- ▶ Placebo (a treatment that appears to be real but contains no medicine)
- ▶ DUPIXENT with a topical corticosteroid (TCS)
- ▶ Placebo with TCS

*Controlled trials have specific protocols that the physician and patient must follow. There are other kinds of clinical trials, but controlled trials are required for FDA approval.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

5 YEAR SAFETY RESULTS IN ADULTS



LONG-TERM SAFETY RESULTS IN ADULTS IN AN EXTENSION STUDY WERE CONSISTENT WITH THOSE SEEN IN THE ORIGINAL CLINICAL TRIALS

The most common side effects observed in clinical trials were: injection site reactions, eye problems, including eye and eyelid inflammation, redness, swelling, itching, eye infection, dry eye, and blurred vision, cold sores in your mouth or on your lips, and high count of a certain white blood cell (eosinophilia).

IMPORTANT CONSIDERATIONS



FDA-APPROVED TO TREAT AGES 6 MONTHS AND UP



STUDIED IN 19 CLINICAL TRIALS: 7 SPECIFICALLY IN ECZEMA, 12 IN 5 OTHER CONDITIONS



NO INITIAL OR ROUTINE BLOOD WORK REQUIRED, PER PRESCRIBING INFORMATION



NOT A STEROID OR IMMUNOSUPPRESSANT



#1 PRESCRIBED BIOLOGIC FOR ECZEMA BY DERMATOLOGISTS AND ALLERGISTS

HEAR FROM ACTUAL PATIENTS

DUPIXENT has helped many of our ambassadors feel the heal and see the difference. [Click below to hear their stories.](#)

Individual results may vary.



ANTHONY,
ACTUAL PATIENT SINCE 2017

“Today I finally feel like I have some control over something that was uncontrolled for too long.”



RACHEL,
ACTUAL PATIENT SINCE 2017

“Being on Dupixent... I worry less about my skin. Eczema is not my first thought when I go outside - and that feels great.”



HEAR FROM MORE PATIENT AMBASSADORS [HERE](#)

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

DUPIXENT HELPS YOU FEEL THE HEAL AND SEE THE DIFFERENCE

Eczema can affect anyone, regardless of age, ethnicity, or gender, and can look different on various skin tones. In some cases, rashes may look red, gray, brown, or purple.

In the adult clinical trials, the efficacy of DUPIXENT has been generally consistent, regardless of race/ethnicity.

ACTUAL DUPIXENT PATIENTS

Individual results may vary.



Actual 4-year-old clinical trial patient.*



Actual, real-world adult patient. Not a clinical trial patient.†



Actual 10-year-old clinical trial patient.*



ADULTS

CHILDREN

CLICK BELOW TO SEE MORE
RESULTS IN THE BEFORE &
AFTER GALLERY

*All patients were prescribed low or medium-potency topical corticosteroids (TCS) based on the clinical trial program.

†Because this was a real-world patient, other factors may have influenced their treatment results (such as use of other treatments).

IMPORTANT SAFETY INFORMATION: Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have AD and asthma and use an asthma medicine.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

NOTICEABLY CLEARER SKIN AND ITCH RELIEF

TRIAL RESULTS: DUPIXENT vs PLACEBO

TWO SEPARATE 4-MONTH CLINICAL TRIALS WITH ADULTS SHOWED:

~4x

SAW CLEAR OR ALMOST CLEAR SKIN

37% vs 9% not on DUPIXENT (placebo)

NEARLY HALF SAW

AT LEAST
75%

CLEARER SKIN

48% vs 13% not on DUPIXENT

MANY ADULTS SAW

AT LEAST
90%

CLEARER SKIN

33% vs 7% not on DUPIXENT

>3x

HAD SIGNIFICANT ITCH REDUCTION

38% vs 11% not on DUPIXENT



ANTHONY,
ACTUAL PATIENT

IMPORTANT SAFETY INFORMATION: Do not change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.



KENNY,
ACTUAL PATIENT

LONG-TERM RESULTS WITH DUPIXENT[®] (dupilumab)

TRIAL RESULTS: DUPIXENT + TOPICAL CORTICOSTEROIDS (TCS) vs TCS ONLY

ADULTS TOOK PART IN A 1-YEAR TRIAL AND SAW:

LONG-LASTING CLEARER SKIN

At 4 months: 39% on DUPIXENT + TCS vs 12% on TCS only

At both 4 months and 1 year: 22% on DUPIXENT + TCS
vs 7% on TCS only

ITCH RELIEF THAT LASTS

At 4 months: 59% on DUPIXENT + TCS vs 20% on TCS only

At 1 year: 51% on DUPIXENT + TCS vs 13% on TCS only

FAST ITCH RELIEF AFTER **FIRST DOSE**

At 2 weeks: 18% on DUPIXENT + TCS vs 8% on TCS only

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including:

- **Allergic reactions.** DUPIXENT can cause allergic reactions, including skin reactions, that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye or painful red or blue bumps under the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

CLEARER SKIN AND LESS ITCH FOR TEENS

TRIAL RESULTS: DUPIXENT vs PLACEBO

TEENS TOOK PART IN A 4-MONTH CLINICAL TRIAL AND SAW:

SIGNIFICANTLY CLEARER SKIN

12x

SAW CLEAR OR ALMOST CLEAR SKIN

24% vs 2% not on DUPIXENT (placebo)

NEARLY HALF SAW

AT LEAST
75%

CLEARER SKIN

42% vs 8% not on DUPIXENT

SOME SAW

AT LEAST
90%

CLEARER SKIN

23% vs 2% not on DUPIXENT

SIGNIFICANT ITCH REDUCTION

~7x

EXPERIENCED ITCH RELIEF

37% vs 5% not on DUPIXENT



OLIVIA,
ACTUAL PATIENT



YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

1-YEAR SAFETY RESULTS:

Long-term safety results in teens continuing on DUPIXENT in the pediatric extension study through 1 year were consistent with the adult clinical trials.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd):

- **Eye problems.** Tell your healthcare provider right away if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision. Your healthcare provider may send you to an ophthalmologist for an eye exam if needed.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

MORE KIDS SAW RELIEF

TRIAL RESULTS: DUPIXENT + TCS vs TCS ONLY

CHILDREN TOOK PART IN A 4-MONTH TRIAL AND SAW:

NOTICEABLY CLEARER SKIN

>2x

SAW CLEAR OR
ALMOST CLEAR SKIN

Under 66 lb: 30% on DUPIXENT + TCS vs 13% on TCS only
66 lb or more: 39% on DUPIXENT+ TCS vs 10% on TCS only

MORE THAN HALF SAW

AT LEAST
75%

CLEARER
SKIN

Under 66 lb: 75% on DUPIXENT + TCS vs 28% on TCS only
66 lb or more: 75% on DUPIXENT + TCS vs 26% on TCS only

CONSIDERABLE ITCH REDUCTION

~4.5x

HAD
LESS ITCH

Under 66 lb: 54% on DUPIXENT + TCS vs 12% on TCS only
66 lb or more: 61% on DUPIXENT + TCS vs 13% on TCS only

Dosing is based on age and weight. Your child's doctor will tell you how much DUPIXENT to give and how often.



YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

1-YEAR SAFETY RESULTS:

Long-term safety results in children continuing on DUPIXENT with or without TCS in the pediatric extension study through 1 year were consistent with the adult clinical trials.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd):

- **Psoriasis.** This can happen in people with atopic dermatitis who receive DUPIXENT. Tell your healthcare provider about any new skin symptoms. Your healthcare provider may send you to a dermatologist for an examination if needed.
- **Joint aches and pain.** Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

EVEN YOUR YOUNGEST CAN FEEL THE HEAL

TRIAL RESULTS: DUPIXENT + TCS vs TCS ONLY

YOUNG CHILDREN TOOK PART IN A 4-MONTH TRIAL AND SAW:

NOTICEABLY CLEARER SKIN

7x

SAW CLEAR OR ALMOST CLEAR SKIN

28% on DUPIXENT + TCS vs 4% on TCS only

MORE THAN HALF SAW

AT LEAST
75%

CLEARER SKIN

53% on DUPIXENT + TCS vs 11% on TCS only

SIGNIFICANT ITCH REDUCTION

>5x

HAD LESS ITCH

48% on DUPIXENT + TCS vs 9% on TCS only



CAMERON,
ACTUAL PATIENT



YOU CAN TAKE DUPIXENT WITH OR WITHOUT A TOPICAL CORTICOSTEROID (TCS)

1-YEAR SAFETY RESULTS:

Long-term safety results in young children continuing on DUPIXENT with or without TCS in the pediatric extension study through 1 year were consistent with the adult clinical trials. In addition, hand-foot-and-mouth disease and warts were reported. In these reported cases, patients continued treatment with DUPIXENT.

IMPORTANT SAFETY INFORMATION: The most common side effects in patients with eczema include injection site reactions; eye problems, including eye and eyelid inflammation, redness, swelling, itching, eye infection, dry eye, and blurred vision; cold sores in your mouth or on your lips; and high count of a certain white blood cell (eosinophilia).

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

DUPIXENT[®] (dupilumab) AND YOU

YOUR DOCTOR DISCUSSION GUIDE

Wherever you are in your treatment journey, or your child's, this guide can help you organize your thoughts to have a productive conversation with your eczema specialist. So get the conversation started about DUPIXENT as an option. Don't pass on an opportunity to help you or your child feel the heal.

ARE YOU A PATIENT OR A CAREGIVER?

- Patient Caregiver

WHAT ARE YOUR OR YOUR LOVED ONE'S TREATMENT GOALS WITH DUPIXENT?

- Clearer skin Itch relief

HOW LONG HAVE YOU OR YOUR LOVED ONE HAD ECZEMA?

- 0-2 years 2-5 years 5-10 years 10+ years

WHAT TREATMENTS HAVE BEEN PRESCRIBED OR USED TO CONTROL YOUR OR YOUR LOVED ONE'S ECZEMA?

- Topical steroids Biologics None
 Phototherapy Over-the-counter products
 Oral steroids Other _____

DESCRIBE YOUR OR YOUR LOVED ONE'S ECZEMA SYMPTOMS.

RASH SEVERITY

- None Mild Moderate Severe

ITCH INTENSITY

- None Mild Moderate Severe

HOW OFTEN DO THESE SYMPTOMS OCCUR?

ITCHING

- Never Rarely Occasionally Frequently

INFLAMED, DISCOLORED SKIN

- Never Rarely Occasionally Frequently

THICKENED, HARDENED PATCHES OF SKIN

- Never Rarely Occasionally Frequently

HOW OFTEN DO FLARE-UPS OCCUR...

- Every few months Monthly Weekly Daily

...AND FOR HOW LONG?

- 1-5 days 5-10 days 10-15 days 15+ days

WHERE ARE YOUR ECZEMA SYMPTOMS?

- Neck Chest Stomach Front of knees
 Face Elbows Feet Back of knees
 Hands Ankles Scalp Other _____

DESCRIBE ANY ADDITIONAL IMPACT LIVING WITH ECZEMA HAS HAD ON YOU OR YOUR LOVED ONE.



FILL OUT YOUR ANSWERS TO SHARE WITH YOUR OWN DOCTOR, OR YOUR CHILD'S, AT THE NEXT APPOINTMENT

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

PARTNERING WITH YOUR HEALTHCARE TEAM



TRY THESE TIPS TO HAVE A PRODUCTIVE DOCTOR VISIT

Having an open conversation with your healthcare team is the first step in working towards a treatment plan that works for you.

- ▶ **DON'T BE AFRAID TO SPEAK UP AND ADVOCATE FOR YOURSELF OR A LOVED ONE**
- ▶ **TAKE PICTURES AND TRACK YOUR (OR YOUR LOVED ONE'S) SYMPTOMS**
- ▶ **PREPARE YOUR QUESTIONS AND TOPICS FOR DISCUSSION WITH YOUR SPECIALISTS**
- ▶ **TAKE NOTES AND BRING THEM TO YOUR HEALTHCARE TEAM VISITS**



SOPHIA,
ACTUAL PATIENT



FIND A SPECIALIST
HEALTHGRADES™ TOOL



CAREGIVER
RESOURCES PAGE

CLICK BELOW TO FIND EACH
OF THE RESOURCES AVAILABLE
ON DUPIXENT.COM

IMPORTANT SAFETY INFORMATION: Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DOSING FOR DUPIXENT[®]

(dupilumab):

	BODY WEIGHT	INITIAL LOADING DOSE	SUBSEQUENT DOSAGE
ADULTS 18+	Any weight	600 mg (two 300-mg injections)	300 mg every 2 weeks
CHILDREN & TEENS 6 TO 17 YEARS OF AGE	132 lb or more	600 mg (two 300-mg injections)	300 mg every 2 weeks
	66 lb to under 132 lb	400 mg (two 200-mg injections)	200 mg every 2 weeks
	33 lb to under 66 lb	600 mg (two 300-mg injections)	300 mg every 4 weeks
	BODY WEIGHT	INITIAL* AND SUBSEQUENT DOSAGE	
YOUNG CHILDREN 6 MONTHS TO 5 YEARS OF AGE	33 lb to under 66 lb	300 mg every 4 weeks	
	11 lb to under 33 lb	200 mg every 4 weeks	

Your doctor will tell you how much **DUPIXENT** to inject and how often.

DUPIXENT IS AVAILABLE AS A PRE-FILLED PEN OR PRE-FILLED SYRINGE.

Your doctor will decide if you or a caregiver can inject DUPIXENT and provide proper training. DUPIXENT is an injection given under the skin (subcutaneous injection). Remember to rotate the injection site every time you administer DUPIXENT.



PRE-FILLED PEN

available in 200 mg and 300 mg doses for patients aged 2+ years



PRE-FILLED SYRINGE

available in 200 mg and 300 mg doses for patients aged 6+ months



CLICK HERE TO WATCH SUPPLEMENTAL TRAINING VIDEOS ON HOW TO ADMINISTER DUPIXENT

*For young children, no initial loading dose is recommended.

DUPIXENT CAN BE GIVEN BY A DOCTOR IN THE OFFICE OR A CAREGIVER AT HOME:

- ▶ In children 6 months to less than 12 years, DUPIXENT should be given by a caregiver
- ▶ In teens 12-17 years, DUPIXENT should be given by or supervised by an adult



LILA,
ACTUAL PATIENT

IMPORTANT SAFETY INFORMATION: Do not try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider. In children 12 years of age and older, it's recommended DUPIXENT be administered by or under supervision of an adult. In children 6 months to less than 12 years of age, DUPIXENT should be given by a caregiver.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

DUPIXENT myway

WITH YOU EVERY STEP OF THE WAY

Once you've been prescribed DUPIXENT, the *DUPIXENT MyWay* patient support program is here to help answer questions you may have about DUPIXENT. Keep in mind your doctor is still the best resource for you if you have any questions about your eczema or your treatment program.

To enroll in *DUPIXENT MyWay* and access helpful resources, call 1-844-DUPIXENT (1-844-387-4936) and select Option 1.



SOPHIA,
ACTUAL PATIENT

YOU'RE NOT ALONE

It's important to know that *DUPIXENT MyWay* helps adults and caregivers of children who are on or about to start DUPIXENT.



Once you're enrolled in the program, and while your insurance benefits are being confirmed for DUPIXENT, you will receive a Welcome Call from your **Case Manager**, who will explain the support available from *DUPIXENT MyWay*. In addition, your **Case Manager** can offer education and support to help you start and stay on track with your treatment.



Your *DUPIXENT MyWay* **Case Manager** can answer questions about DUPIXENT, arrange supplemental injection training, and help you start and stay on track.



With the *DUPIXENT MyWay* **Patient App**, people just like you can get access to tools to help you start and stay on track with your treatment. Contact your **Case Manager** for an invitation with directions on how to get the app.



The *DUPIXENT MyWay* **Mentor Program** gives you the opportunity to speak with other DUPIXENT patients and caregivers to hear their experiences.



[CLICK HERE TO LEARN ABOUT ADDITIONAL
DUPIXENT MyWay RESOURCES](#)

Please see Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).



YOUR COPAY COULD BE AS LITTLE AS \$0*

The DUPIXENT MyWay Copay Card Program may help reduce the out-of-pocket cost of DUPIXENT. Eligible patients with commercial health insurance may pay as little as \$0* in copay per fill of DUPIXENT® (dupilumab). Terms and conditions apply.

YOU MAY BE ELIGIBLE IF YOU:

- ▶ Have commercial insurance
- ▶ Have a DUPIXENT prescription for an FDA-approved condition
- ▶ Reside in the 50 United States, the District of Columbia, Puerto Rico, Guam, or the USVI
- ▶ Are a patient or caregiver 18 years or older

**Subject to the program maximum per patient per calendar year. Approval is not guaranteed. THIS IS NOT INSURANCE. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE, or other federal or state programs, including any state pharmaceutical assistance programs. This program is not valid where prohibited by law, taxed, or restricted. DUPIXENT MyWay reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. Any savings provided by the program may vary depending on patients' out-of-pocket costs. The program is intended to help patients afford DUPIXENT. Patients may have insurance plans that attempt to dilute the impact of the assistance available under the program. In those situations, the program may change its terms. Additional terms and conditions apply.*

OTHER OPTIONS FOR SUPPORT

Through the Patient Assistance Program, eligible patients who are uninsured or rendered uninsured by their insurance provider could receive DUPIXENT free of charge if they qualify.

Those who may qualify must be at least 18 years of age or older, a resident of the 50 United States, the District of Columbia, Puerto Rico, Guam, or the USVI, and demonstrate a financial need.

Call DUPIXENT MyWay® at 1-844-387-4936 for full program information and eligibility criteria.



TYLER,
ACTUAL PATIENT

DUPIXENT[®]
(dupilumab) Injection
200mg · 300mg

OVER 645,000* PATIENTS
PRESCRIBED DUPIXENT

HELP
HEAL
YOUR
SKIN
FROM
WITHIN[™]

LILA,
ACTUAL PATIENT



CLICK HERE TO
LEARN MORE

*New adult and pediatric patients (6 months of age and older) with uncontrolled moderate-to-severe eczema. IQVIA National Source of Business (NSOB) data through March 2025.

ASK YOUR ECZEMA SPECIALIST ABOUT DUPIXENT
AND VISIT [DUPIXENT.COM](https://www.dupixent.com) OR CALL 1-844-DUPIXENT (1-844-387-4936)

IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT[®].

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

sanofi | REGENERON[®]

© 2026 Sanofi and Regeneron Pharmaceuticals, Inc. All rights reserved.
DUPIXENT[®] and DUPIXENT MyWay[®] are registered trademarks of
Sanofi Biotechnology. US.DUP.26.04.0469 04/26